

## UNKL rabbit pAb

<b>Catalog No :</b>	YT7712
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB
<b>Target :</b>	UNKL
<b>Gene Name :</b>	UNKL C16orf28 ZC3H5L ZC3HDC5L
<b>Protein Name :</b>	UNKL
<b>Human Gene Id :</b>	64718
<b>Human Swiss Prot No :</b>	Q9H9P5
<b>Mouse Gene Id :</b>	74154
<b>Mouse Swiss Prot No :</b>	Q5FWH2
<b>Immunogen :</b>	Synthesized peptide derived from human UNKL AA range: 296-346
<b>Specificity :</b>	This antibody detects endogenous levels of UNKL at Human/Mouse
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1[?]500-2000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)


**Molecularweight :** 75kD

**Background :** This gene encodes a RING finger protein that may function in Rac signaling. It can bind to Brg/Brm-associated factor 60b and can promote its ubiquitination. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jun 2013],

**Function :** similarity:Belongs to the unkempt family.,similarity:Contains 1 RING-type zinc finger.,

**Subcellular Location :** [Isoform 4]: Cytoplasm. Nucleus. Isoform 4 is primarily localized in the cytoplasm but has the ability to shuttle between the nucleus and the cytoplasm.

## Products Images



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4° over night