

MDSP rabbit pAb

Catalog No :	YT7741
Reactivity :	Human;Mouse
Applications :	WB
Target :	MDSP
Gene Name :	DUSP13 BEDP MDSP
Protein Name :	MDSP
Human Gene Id :	51207
Human Swiss Prot	Q6B8I1
NO : Mouse Gene Id :	27389
Mouse Swiss Prot	Q6B8I0
NO : Immunogen :	Synthesized peptide derived from human MDSP AA range: 48-98
Specificity :	This antibody detects endogenous levels of MDSP at Human/Mouse
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1 [9500-2000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)



Molecularweight: 21kD

Background :	Members of the protein-tyrosine phosphatase superfamily cooperate with protein kinases to regulate cell proliferation and differentiation. This superfamily is separated into two families based on the substrate that is dephosphorylated. One family, the dual specificity phosphatases (DSPs) acts on both phosphotyrosine and phosphoserine/threonine residues. This gene encodes different but related DSP proteins through the use of non-overlapping open reading frames, alternate splicing, and presumed different transcription promoters. Expression of the distinct proteins from this gene has been found to be tissue specific and the proteins may be involved in postnatal development of specific tissues. A protein encoded by the upstream ORF was found in skeletal muscle, whereas the encoded protein from the downstream ORF was found only in testis. In mouse, a similar pattern of expression was found. Multiple alternatively spliced transcript variants were described, but the full-length sequence of only some were determined. [provided by RefSeq, Jul 2008],
Function :	catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate.,catalytic activity:Protein tyrosine phosphate + H(2)O = protein tyrosine + phosphate.,function:May be involved in the regulation of meiosis and/or differentiation of testicular germ cells during spermatogenesis. Exhibits intrinsic phosphatase activity towards both phospho-seryl/threonyl and -tyrosyl residues of myelin basic protein, with similar specific activities in vitro.,similarity:Belongs to the protein-tyrosine phosphatase family. Non-receptor class dual specificity subfamily.,similarity:Contains 1 tyrosine-protein phosphatase domain.,tissue specificity:Most abundantly expressed in the testis.,
Subcellular	Cytoplasm .
Location :	Skeletal muscle specific
LAPIC331011.	

