

ODO2 rabbit pAb

Catalog No :	YT8137
Reactivity :	Human;Mouse;Rat
Applications :	IHC;WB
Target :	DLST
Gene Name :	DLST DLTS
Protein Name :	Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial (EC 2.3.1.61) (2-oxoglutarate dehydrogenase complex component E2) (OGDC-E2) (Dihydrolip
Human Gene Id :	1743
Human Swiss Prot No :	P36957
Mouse Gene Id :	78920
Mouse Swiss Prot No :	Q9D2G2
Rat Gene Id :	299201
Rat Swiss Prot No :	Q01205
Immunogen :	Synthesized peptide derived from human C-terminal ODO2
Specificity :	This antibody detects endogenous levels of ODO2 at Human, Mouse,Rat
Formulation :	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000 IHC 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight : 50kD

Function : Dihydrolipoamide succinyltransferase (E2) component of the 2-oxoglutarate dehydrogenase complex. The 2-oxoglutarate dehydrogenase complex catalyzes the overall conversion of 2-oxoglutarate to succinyl-CoA and CO₂. The 2-oxoglutarate dehydrogenase complex is mainly active in the mitochondrion . A fraction of the 2-oxoglutarate dehydrogenase complex also localizes in the nucleus and is required for lysine succinylation of histones: associates with KAT2A on chromatin and provides succinyl-CoA to histone succinyltransferase KAT2A .

Subcellular Location : Mitochondrion matrix . Nucleus . Mainly localizes in the mitochondrion. A small fraction localizes to the nucleus, where the 2-oxoglutarate dehydrogenase complex is required for histone succinylation. .

Products Images