

## Cleaved-Cathepsin D LC (G65) Polyclonal Antibody

<b>Catalog No :</b>	YC0038
<b>Reactivity :</b>	Human;Monkey
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	Cathepsin D
<b>Fields :</b>	>>Sphingolipid signaling pathway;>>Autophagy - animal;>>Lysosome;>>Apoptosis;>>Estrogen signaling pathway;>>Tuberculosis;>>Diabetic cardiomyopathy
<b>Gene Name :</b>	CTSD
<b>Protein Name :</b>	Cathepsin D
<b>Human Gene Id :</b>	1509
<b>Human Swiss Prot No :</b>	P07339
<b>Mouse Swiss Prot No :</b>	P18242
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human CATD. AA range:46-95
<b>Specificity :</b>	Cleaved-Cathepsin D LC (G65) Polyclonal Antibody detects endogenous levels of fragment of activated Cathepsin D LC protein resulting from cleavage adjacent to G65.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. ELISA: 1:20000. Not yet tested in other applications.
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

**Concentration :** 1 mg/ml

---

**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)

---

**Observed Band :** 17kD

---

**Cell Pathway :** Lysosome;

---

**Background :** This gene encodes a member of the A1 family of peptidases. The encoded preproprotein is proteolytically processed to generate multiple protein products. These products include the cathepsin D light and heavy chains, which heterodimerize to form the mature enzyme. This enzyme exhibits pepsin-like activity and plays a role in protein turnover and in the proteolytic activation of hormones and growth factors. Mutations in this gene play a causal role in neuronal ceroid lipofuscinosis-10 and may be involved in the pathogenesis of several other diseases, including breast cancer and possibly Alzheimer's disease. [provided by RefSeq, Nov 2015],

---

**Function :** catalytic activity:Specificity similar to, but narrower than, that of pepsin A. Does not cleave the 4-Gln-|-His-5 bond in B chain of insulin.,disease:Defects in CTSD are the cause of neuronal ceroid lipofuscinosis 10 (CLN10) [MIM:610127]; also known as neuronal ceroid lipofuscinosis due to cathepsin D deficiency. The neuronal ceroid lipofuscinosis are a group of progressive neurodegenerative diseases in children and in adults, characterized by visual and mental decline, motor disturbance, epilepsy and behavioral changes.,function:Acid protease active in intracellular protein breakdown. Involved in the pathogenesis of several diseases such as breast cancer and possibly Alzheimer disease.,polymorphism:The Val-58 allele is significantly overrepresented in demented patients (11.8%) compared with non-demented controls (4.9%). Carriers of the Val-58 allele have a 3.1-fold increased risk for de

---


**Subcellular Location :** Lysosome. Melanosome. Secreted, extracellular space. Identified by mass spectrometry in melanosome fractions from stage I to stage IV. In aortic samples, detected as an extracellular protein loosely bound to the matrix (PubMed:20551380).

---

**Expression :** Expressed in the aorta extracellular space (at protein level) (PubMed:20551380). Expressed in liver (at protein level) (PubMed:1426530).

---

## Products Images



Western Blot analysis of various cells using Cleaved-Cathepsin D LC (G65) Polyclonal Antibody