

ENTK (light chain, Cleaved-Ile785) rabbit pAb

Catalog No: YC0153

**Reactivity:** Human; Rat; Mouse;

**Applications:** WB;ELISA

Target: ENTK

Gene Name: TMPRSS15 ENTK PRSS7

Protein Name: ENTK (light chain, Cleaved-Ile785)

P98073

P97435

Human Gene Id: 5651

**Human Swiss Prot** 

No:

Mouse Gene ld: 19146

**Mouse Swiss Prot** 

No:

Immunogen: Synthesized peptide derived from human ENTK (light chain, Cleaved-Ile785)

**Specificity:** This antibody detects endogenous levels of Human ENTK (light chain, Cleaved-

lle785, protein was cleaved amino acid sequence between 785-786)

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:1000-2000 ELISA 1:5000-20000

**Purification:** The antibody was affinity-purified from rabbit serum by affinity-chromatography

using specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/2



Observed Band: 26kD

**Background:** catalytic activity:Activation of trypsinogen by selective cleavage of 6-Lys-|-lle-7

bond., disease: Defects in PRSS7 are a cause of enterokinase deficiency

[MIM:226200]; a life-threatening intestinal malabsorption disorder characterized by diarrhea and failure to thrive.,function:Responsible for initiating activation of pancreatic proteolytic proenzymes (trypsin, chymotrypsin and carboxypeptidase A). It catalyzes the conversion of trypsinogen to trypsin which in turn activates other proenzymes including chymotrypsinogen, procarboxypeptidases, and proelastases..PTM:The chains are derived from a single precursor that is cleaved

by a trypsin-like protease., similarity: Belongs to the peptidase S1

family.,similarity:Contains 1 MAM domain.,similarity:Contains 1 peptidase S1

domain.,similarity:Contains 1 SEA domain.,similarity:Contains 1 SRCR domain.,similarity:Contains 2 CUB domains.,similarity:Contains 2 LDL-receptor

class A domains., subunit: Heterodimer of a catalytic (light) chain and a

multidomain (heavy) chain linked by a disulfide bond.,tissue specificity:Intestinal

brush border.,

Function: proteolysis,

Subcellular Location:

Membrane; Single-pass type II membrane protein.

**Expression:** Intestinal brush border.

## **Products Images**

2/2