

**CD19 (phospho Tyr531) Polyclonal Antibody**

<b>Catalog No :</b>	YP0295
<b>Reactivity :</b>	Human;Mouse;Monkey
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	CD19
<b>Fields :</b>	>>PI3K-Akt signaling pathway;>>Hematopoietic cell lineage;>>B cell receptor signaling pathway;>>Epstein-Barr virus infection;>>Primary immunodeficiency
<b>Gene Name :</b>	CD19
<b>Protein Name :</b>	B-lymphocyte antigen CD19
<b>Human Gene Id :</b>	930
<b>Human Swiss Prot No :</b>	P15391
<b>Mouse Gene Id :</b>	12478
<b>Mouse Swiss Prot No :</b>	P25918
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human CD19 around the phosphorylation site of Tyr531. AA range:501-550
<b>Specificity :</b>	Phospho-CD19 (Y531) Polyclonal Antibody detects endogenous levels of CD19 protein only when phosphorylated at Y531.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

**Concentration :** 1 mg/ml

---

**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)

---

**Observed Band :** 61kD

---

**Cell Pathway :** Hematopoietic cell lineage;B\_Cell\_Antigen;Primary immunodeficiency;

---

**Background :** Lymphocytes proliferate and differentiate in response to various concentrations of different antigens. The ability of the B cell to respond in a specific, yet sensitive manner to the various antigens is achieved with the use of low-affinity antigen receptors. This gene encodes a cell surface molecule which assembles with the antigen receptor of B lymphocytes in order to decrease the threshold for antigen receptor-dependent stimulation. [provided by RefSeq, Jul 2008],

---

**Function :** disease:Defects in CD19 are a cause of hypogammaglobulinemia [MIM:107265].,function:Assembles with the antigen receptor of B lymphocytes in order to decrease the threshold for antigen receptor-dependent stimulation.,online information:CD19 mutation db,PTM:Phosphorylated on serine and threonine upon DNA damage, probably by ATM or ATR. Phosphorylated on tyrosine following B-cell activation.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Forms a complex with CD21, CD81 and CD225 in the membrane of mature B cells. Interacts with VAV. Interacts with GRB2 and SOS when phosphorylated on Tyr-348 and/or Tyr-378. Interacts with PLCG2 when phosphorylated on Tyr-409.,

---


**Subcellular Location :** Cell membrane ; Single-pass type I membrane protein . Membrane raft ; Single-pass type I membrane protein .

---

**Expression :** Detected on marginal zone and germinal center B cells in lymph nodes (PubMed:2463100). Detected on blood B cells (at protein level) (PubMed:2463100, PubMed:16672701).

---

## Products Images



Western blot analysis of lysates from COS7 cells treated with Serum 10% 30', using CD19 (Phospho-Tyr531) Antibody. The lane on the right is blocked with the phospho peptide.