

PLC β 3 (phospho Ser1105) Polyclonal Antibody

Catalog No :	YP0606
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	PLC β 3
Fields :	>>Inositol phosphate metabolism;>>Metabolic pathways;>>Rap1 signaling pathway;>>Calcium signaling pathway;>>cGMP-PKG signaling pathway;>>Chemokine signaling pathway;>>Phosphatidylinositol signaling system;>>Sphingolipid signaling pathway;>>Phospholipase D signaling pathway;>>Adrenergic signaling in cardiomyocytes;>>Vascular smooth muscle contraction;>>Wnt signaling pathway;>>Apelin signaling pathway;>>Gap junction;>>Platelet activation;>>Neutrophil extracellular trap formation;>>NOD-like receptor signaling pathway;>>Circadian entrainment;>>Long-term potentiation;>>Retrograde endocannabinoid signaling;>>Glutamatergic synapse;>>Cholinergic synapse;>>Serotonergic synapse;>>Dopaminergic synapse;>>Long-term depression;>>Taste transduction;>>Inflammatory mediator regulation of TRP channels;>>Insulin secretion;>>GnRH signaling pathway;>>Estrogen signaling pathway;>>Melanogenesis;>>Thyroid hormone synthesis;>>Thyroid hormone signaling pathway;>>Oxytocin signaling pathway;>>Glucagon signaling p
Gene Name :	PLCB3
Protein Name :	1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase beta-3
Human Gene Id :	5331
Human Swiss Prot No :	Q01970
Mouse Swiss Prot No :	P51432
Rat Swiss Prot No :	Q99JE6
Immunogen :	The antiserum was produced against synthesized peptide derived from human PLCB3 around the phosphorylation site of Ser1105. AA range:1071-1120
Specificity :	Phospho-PLC β 3 (S1105) Polyclonal Antibody detects endogenous levels of

PLC β 3 protein only when phosphorylated at S1105.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source : Polyclonal, Rabbit,IgG

Dilution : WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000.. IF 1:50-200

Purification : The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 160kD

Cell Pathway : Stem cell pathway; WNT;WNT-T CELL; β -Catenin; AMPK


Background : This gene encodes a member of the phosphoinositide phospholipase C beta enzyme family that catalyze the production of the secondary messengers diacylglycerol and inositol 1,4,5-triphosphate from phosphatidylinositol in G-protein-linked receptor-mediated signal transduction. Alternative splicing results in multiple transcript variants.[provided by RefSeq, May 2010],

Function : catalytic activity:1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate + H(2)O = 1D-myo-inositol 1,4,5-trisphosphate + diacylglycerol.,cofactor:Calcium.,function:The production of the second messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes.,similarity:Contains 1 C2 domain.,similarity:Contains 1 PI-PLC X-box domain.,similarity:Contains 1 PI-PLC Y-box domain.,subunit:Interacts with SHANK2 (By similarity). Interacts with LPAR2.,


Subcellular Location : Cytoplasm . Membrane . Nucleus . And particulate fractions. .

Expression : Epithelium,Uterus,


Products Images




Western Blot analysis of various cells using Phospho-PLC β 3 (S1105) Polyclonal Antibody diluted at 1:1000




Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100 (4° overnight). High-pressure and temperature Tris-EDTA, pH 8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using PLCB3 (Phospho-Ser1105) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using PLCB3 (Phospho-Ser1105) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from A431 cells, using PLCB3 (Phospho-Ser1105) Antibody. The lane on the right is blocked with the phospho peptide.