

**BLNK (phospho Tyr96) Polyclonal Antibody**

<b>Catalog No :</b>	YP0687
<b>Reactivity :</b>	Human;Mouse;Monkey
<b>Applications :</b>	WB;IHC;IF;ELISA
<b>Target :</b>	BLNK
<b>Fields :</b>	>>NF-kappa B signaling pathway;>>Osteoclast differentiation;>>B cell receptor signaling pathway;>>Epstein-Barr virus infection;>>Primary immunodeficiency
<b>Gene Name :</b>	BLNK
<b>Protein Name :</b>	B-cell linker protein
<b>Human Gene Id :</b>	29760
<b>Human Swiss Prot No :</b>	Q8WV28
<b>Mouse Gene Id :</b>	17060
<b>Mouse Swiss Prot No :</b>	Q9QUN3
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human BLNK around the phosphorylation site of Tyr96. AA range:62-111
<b>Specificity :</b>	Phospho-BLNK (Y96) Polyclonal Antibody detects endogenous levels of BLNK protein only when phosphorylated at Y96.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000.. IF 1:50-200
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

**Concentration :** 1 mg/ml

**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)

**Observed Band :** 50kD

**Cell Pathway :** B\_Cell\_Antigen;Primary immunodeficiency;

**Background :** This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell development. This protein bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. The phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct signaling effectors following B cell receptor activation. Mutations in this gene cause hypoglobulinemia and absent B cells, a disease in which the pro- to pre-B-cell transition is developmentally blocked. Deficiency in this protein has also been shown in some cases of pre-B acute lymphoblastic leukemia. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, May 2012],


**Function :** disease:Defects in BLNK are the cause of hypoglobulinemia and absent B-cells [MIM:604515]. This is a developmental blockage at the pro- to pre-B-cell transition.,disease:In 6 of 34 childhood pre-B acute lymphoblastic leukemia (ALL) samples that were tested showed a complete loss or drastic reduction of BLNK expression.,function:Functions as a central linker protein that bridges kinases associated with the B-cell receptor (BCR) with a multitude of signaling pathways, regulating biological outcomes of B-cell function and development. Plays a role in the activation of ERK/EPHB2, MAP kinase p38 and JNK. Modulates AP1 activation. Important for the activation of NF-kappa-B and NFAT. Plays an important role in BCR-mediated PLCG1 and PLCG2 activation and Ca(2+) mobilization and is required for trafficking of the BCR to late endosomes. However, does not seem to be required for pre-BCR-mediated ac

**Subcellular Location :** Cytoplasm . Cell membrane . BCR activation results in the translocation to membrane fraction.


**Expression :** Expressed in B-cell lineage and fibroblast cell lines (at protein level). Highest levels of expression in the spleen, with lower levels in the liver, kidney, pancreas, small intestines and colon.

---


## Products Images




Western Blot analysis of COLO205 cells using Phospho-BLNK (Y96) Polyclonal Antibody



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using BLNK (Phospho-Tyr96) Antibody



Immunohistochemistry analysis of paraffin-embedded human lymph node, using BLNK (Phospho-Tyr96) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from COLO205 cells, using BLNK (Phospho-Tyr96) Antibody. The lane on the right is blocked with the phospho peptide.