


## MEK Kinase-1 (phospho Thr1402) Polyclonal Antibody


<b>Catalog No :</b>	YP0786
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;IHC;IF;ELISA
<b>Target :</b>	MEK Kinase-1
<b>Fields :</b>	>>MAPK signaling pathway;>>Ubiquitin mediated proteolysis;>>Tight junction;>>RIG-I-like receptor signaling pathway;>>Neurotrophin signaling pathway;>>GnRH signaling pathway;>>Growth hormone synthesis, secretion and action;>>Hepatitis B;>>Human T-cell leukemia virus 1 infection
<b>Gene Name :</b>	MAP3K1
<b>Protein Name :</b>	Mitogen-activated protein kinase kinase kinase 1
<b>Human Gene Id :</b>	4214
<b>Human Swiss Prot No :</b>	Q13233
<b>Mouse Swiss Prot No :</b>	P53349
<b>Rat Gene Id :</b>	116667
<b>Rat Swiss Prot No :</b>	Q62925
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human MAP3K1 around the phosphorylation site of Thr1402. AA range:1368-1417
<b>Specificity :</b>	Phospho-MEK Kinase-1 (T1402) Polyclonal Antibody detects endogenous levels of MEK Kinase-1 protein only when phosphorylated at T1402.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200

<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	160kD
<b>Cell Pathway :</b>	SAPK_JNK; Regulation of Actin Dynamics; Cell Growth; Stem cell pathway; MAPK_ERK_Growth;MAPK_G_Protein; B Cell Receptor
<b>Background :</b>	The protein encoded by this gene is a serine/threonine kinase and is part of some signal transduction cascades, including the ERK and JNK kinase pathways as well as the NF-kappa-B pathway. The encoded protein is activated by autophosphorylation and requires magnesium as a cofactor in phosphorylating other proteins. This protein has E3 ligase activity conferred by a plant homeodomain (PHD) in its N-terminus and phospho-kinase activity conferred by a kinase domain in its C-terminus. [provided by RefSeq, Mar 2012],
<b>Function :</b>	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by autophosphorylation on Thr-1400 and Thr-1412 following oligomerization.,function:Component of a protein kinase signal transduction cascade. Activates the ERK and JNK kinase pathways by phosphorylation of MAP2K1 and MAP2K4. Activates CHUK and IKBKB, the central protein kinases of the NF-kappa-B pathway.,PTM:Autophosphorylated.,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase kinase subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 RING-type zinc finger.,similarity:Contains 1 SWIM-type zinc finger.,subunit:Binds both upstream activators and downstream substrates in multimolecular complexes through its N-terminus. Oligomerizes after binding MAP4K2 or TRAF2. Interacts with AXIN1.,
<b>Subcellular Location :</b>	cytoplasm,cytosol,
<b>Expression :</b>	Leukocyte,


## Products Images




Western Blot analysis of NIH-3T3 cells using Phospho-MEK Kinase-1 (T1402) Polyclonal Antibody diluted at 1:2000



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using MAP3K1 (Phospho-Thr1402) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using MAP3K1 (Phospho-Thr1402) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from Jurkat cells, using MAP3K1 (Phospho-Thr1402) Antibody. The lane on the right is blocked with the phospho peptide.