

SREBP-1 (phospho Ser439) Polyclonal Antibody

Catalog No: YP0833

Reactivity: Human;Rat

Applications: WB;IHC;IF;ELISA

Target: SREBP-1

Fields: >>AMPK signaling pathway;>>Insulin signaling pathway;>>Insulin

resistance;>>Non-alcoholic fatty liver disease;>>Alcoholic liver disease

Gene Name: SREBF1

Protein Name: Sterol regulatory element-binding protein 1

P36956

Q9WTN3

Human Gene Id: 6720

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Rat Gene Id: 78968

Rat Swiss Prot No: P56720

Immunogen: The antiserum was produced against synthesized peptide derived from human

SREBP-1 around the phosphorylation site of Ser439. AA range:405-454

Specificity: Phospho-SREBP-1 (S439) Polyclonal Antibody detects endogenous levels of

SREBP-1 protein only when phosphorylated at S439.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, lgG

Dilution: WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:5000.. IF 1:50-200

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

1/3



chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 122kD

Cell Pathway: AMPK; Protein_Acetylation

Background: This gene encodes a transcription factor that binds to the sterol regulatory

element-1 (SRE1), which is a decamer flanking the low density lipoprotein receptor gene and some genes involved in sterol biosynthesis. The protein is synthesized as a precursor that is attached to the nuclear membrane and endoplasmic reticulum. Following cleavage, the mature protein translocates to the nucleus and activates transcription by binding to the SRE1. Sterols inhibit the cleavage of the precursor, and the mature nuclear form is rapidly catabolized, thereby reducing transcription. The protein is a member of the basic helix-loophelix-leucine zipper (bHLH-Zip) transcription factor family. This gene is located within the Smith-Magenis syndrome region on chromosome 17. [provided by

RefSeq, Mar 2016],

Function: alternative products:Additional isoforms seem to exist, function: Transcriptional

activator required for lipid homeostasis. Regulates transcription of the LDL receptor gene as well as the fatty acid and to a lesser degree the cholesterol synthesis pathway (By similarity). Binds to the sterol regulatory element 1 (SRE-1) (5'-ATCACCCCAC-3'). Has dual sequence specificity binding to both an E-box motif (5'-ATCACGTGA-3') and to SRE-1 (5'-ATCACCCCAC-3').,online

cholesterol the SCAP/SREBP complex is recruited into COPII vesicles for export from the ER. In the Golgi complex SREBPs are cleaved sequentially by site-1 and site-2 protease. The first cleavage by site-1 protease occurs within the luminal

loop, the second cleavage by site-2 protease occurs within the first

information: Sterol regulatory element-binding protein entry, PTM: At low

transmembrane domain and releases the transcription factor fr

Subcellular [Sterol regulatory element-binding protein 1]: Endoplasmic reticulum membrane ; **Location :** Multi-pass membrane protein . Golgi apparatus membrane ; Multi-pass

Multi-pass membrane protein . Golgi apparatus membrane ; Multi-pass membrane protein . Cytoplasmic vesicle, COPII-coated vesicle membrane ; Multi-

pass membrane protein . At high sterol concentrations, the SCAP-SREBP is retained in the endoplasmic reticulum. Low sterol concentrations promote recruitment into COPII-coated vesicles and transport of the SCAP-SREBP to the

Golgi, where it is processed. .; [Processed sterol regulatory element-binding protein 1]: Nucleus .; [Isoform SREBP-1aDelta]: Nucleus .; [Isoform

SREBP-1cDelta]: Nucleus.

Expression: Expressed in a wide variety of tissues, most abundant in liver and adrenal gland


(PubMed:8402897). In fetal tissues lung and liver shows highest expression

2/3




(PubMed:8402897).; [Isoform SREBP-1A]: Predominates in hepatoma cell lines (PubMed:8402897). Also expressed in kidney, brain, white fat, and muscle (PubMed:8402897).; [Isoform SREBP-1C]: Predominantly expressed in liver and adipose tissues (PubMed:8402897). Also expressed in kidney, brain, white fat, and muscle (PubMed:8402897).

Products Images



Immunohistochemistry analysis of paraffin-embedded human brain, using SREBP-1 (Phospho-Ser439) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from Jurkat cells treated with TNF 20ng/ml 30', using SREBP-1 (Phospho-Ser439) Antibody. The lane on the right is blocked with the phospho peptide.