


**KIR3.1 (phospho Ser185) Polyclonal Antibody**

<b>Catalog No :</b>	YP0932
<b>Reactivity :</b>	Human;Mouse;Rat;Monkey
<b>Applications :</b>	WB;IHC;IF;ELISA
<b>Target :</b>	KIR3.1
<b>Fields :</b>	>>Circadian entrainment;>>Retrograde endocannabinoid signaling;>>Glutamatergic synapse;>>Cholinergic synapse;>>Serotonergic synapse;>>Dopaminergic synapse;>>Estrogen signaling pathway;>>Oxytocin signaling pathway;>>GnRH secretion;>>Morphine addiction
<b>Gene Name :</b>	KCNJ3
<b>Protein Name :</b>	G protein-activated inward rectifier potassium channel 1
<b>Human Gene Id :</b>	3760
<b>Human Swiss Prot No :</b>	P48549
<b>Mouse Gene Id :</b>	16519
<b>Mouse Swiss Prot No :</b>	P63250
<b>Rat Gene Id :</b>	50599
<b>Rat Swiss Prot No :</b>	P63251
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human GIRK1/KIR3.1/KCNJ3 around the phosphorylation site of Ser185. AA range:151-200
<b>Specificity :</b>	Phospho-KIR3.1 (S185) Polyclonal Antibody detects endogenous levels of KIR3.1 protein only when phosphorylated at S185.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.


<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	50kD
<b>Background :</b>	<p>Potassium channels are present in most mammalian cells, where they participate in a wide range of physiologic responses. The protein encoded by this gene is an integral membrane protein and inward-rectifier type potassium channel. The encoded protein, which has a greater tendency to allow potassium to flow into a cell rather than out of a cell, is controlled by G-proteins and plays an important role in regulating heartbeat. It associates with three other G-protein-activated potassium channels to form a heteromultimeric pore-forming complex that also couples to neurotransmitter receptors in the brain and whereby channel activation can inhibit action potential firing by hyperpolarizing the plasma membrane. These multimeric G-protein-gated inwardly-rectifying potassium (GIRK) channels may play a role in the pathophysiology of epilepsy, addiction, Down's syndrome, at</p>
<b>Function :</b>	<p>function:This potassium channel is controlled by G proteins. Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium. This receptor plays a crucial role in regulating the heartbeat.,similarity:Belongs to the inward rectifier-type potassium channel family.,subunit:Associates with GIRK2, GIRK3 or GIRK4 to form a G-protein activated heteromultimer pore-forming unit. The resulting inward current is much larger.,</p>
<b>Subcellular Location :</b>	Membrane; Multi-pass membrane protein.
<b>Expression :</b>	Brain,Epithelium,

---


## Products Images




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using GIRK1/KIR3.1/KCNJ3 (Phospho-Ser185) Antibody



Immunofluorescence analysis of HeLa cells, using GIRK1/KIR3.1/KCNJ3 (Phospho-Ser185) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma, using GIRK1/KIR3.1/KCNJ3 (Phospho-Ser185) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from RAW264.7 cells treated with Insulin 0.01U/ml 15', using GIRK1/KIR3.1/KCNJ3 (Phospho-Ser185) Antibody. The lane on the right is blocked with the phospho peptide.