

PA24B Polyclonal Antibody

| | |
|------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Catalog No : | YN0632 |
| Reactivity : | Human;Mouse |
| Applications : | WB;ELISA |
| Target : | PA24B |
| Fields : | >>Glycerophospholipid metabolism;>>Ether lipid metabolism;>>Arachidonic acid metabolism;>>Linoleic acid metabolism;>>alpha-Linolenic acid metabolism;>>Metabolic pathways;>>MAPK signaling pathway;>>Ras signaling pathway;>>Phospholipase D signaling pathway;>>Necroptosis;>>Vascular smooth muscle contraction;>>VEGF signaling pathway;>>Platelet activation;>>Fc epsilon RI signaling pathway;>>Fc gamma R-mediated phagocytosis;>>Glutamatergic synapse;>>Serotonergic synapse;>>Long-term depression;>>Inflammatory mediator regulation of TRP channels;>>GnRH signaling pathway;>>Ovarian steroidogenesis;>>Oxytocin signaling pathway;>>Choline metabolism in cancer |
| Gene Name : | PLA2G4B |
| Protein Name : | Cytosolic phospholipase A2 beta (cPLA2-beta) (EC 3.1.1.4) (Phospholipase A2 group IVB) |
| Human Gene Id : | 100137049 |
| Human Swiss Prot No : | P0C869 |
| Mouse Swiss Prot No : | P0C871 |
| Immunogen : | Synthesized peptide derived from part region of human protein |
| Specificity : | PA24B Polyclonal Antibody detects endogenous levels of protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | WB 1:500-2000 ELISA 1:5000-20000 |

| | |
|-------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |
| Observed Band : | 85kD |
| Cell Pathway : | Glycerophospholipid metabolism;Ether lipid metabolism;Arachidonic acid metabolism;Linoleic acid metabolism;alpha-Linolenic acid metabolism;MAPK_ERK_Growth;MAPK_G_Protein;Vascular smooth muscle contrac |
| Background : | This locus represents naturally-occurring readthrough transcription between the neighboring jumonji domain containing 7 (JMJD7) and phospholipase A2, group IVB (cytosolic) (PLA2G4B) genes. Readthrough transcripts encode fusion proteins that share amino acid sequence with each individual gene product, including a partial JmjC domain and downstream C2 and phospholipase A2 domains. Alternatively spliced transcript variants have been observed. [provided by RefSeq, Oct 2013], |
| Function : | catalytic activity:Phosphatidylcholine + H(2)O = 1-acylglycerophosphocholine + a carboxylate.,caution:Most tissues also express read-through transcripts from this gene into the upstream gene (Jmjd7), some of which may encode fusion proteins.,caution:This sequence was first thought to be an alternatively spliced isoform of Pla2g4b. It is derived from Jmjd7 which is located upstream of Pla2g4b. Most tissues also express read-through transcripts from Jmjd7 into the downstream Pla2g4b gene, some of which may encode fusion proteins combining the N-terminus of this protein with PLA2G4B protein.,domain:The N-terminal C2 domain associates with lipid membranes and mediates its regulation by presenting the active site to its substrate in response to elevations of cytosolic Ca(2+).,enzyme regulation:Stimulated by cytosolic Ca(2+).,function:Calcium-dependent phospholipase A2 that selectively hydroly |
| Subcellular Location : | [Isoform 3]: Cytoplasm, cytosol . Mitochondrion membrane ; Peripheral membrane protein. Early endosome membrane ; Peripheral membrane protein. Translocates to membrane vesicles in a calcium-dependent fashion. . ; [Isoform 5]: Cytoplasm, cytosol . |
| Expression : | Widely expressed. Expressed at higher level in brain, heart, liver, cerebellum and pancreas. |

Products Images