

EPHA1 (Phospho Tyr605) Polyclonal Antibody

Catalog No: YP1229

Reactivity: Human; Mouse

Applications: IHC;IF;WB

Target: EPHA1

Fields: >>Axon guidance

Gene Name: EPHA1 EPH EPHT EPHT1

Protein Name: EPHA1 (Phospho-Tyr605)

P21709

Human Gene Id: 2041

Human Swiss Prot

No:

Immunogen: Synthesized peptide derived from human EPHA1 (Phospho-Tyr605)

Specificity: This antibody detects endogenous phospho levels of EPHA1 (Phospho-Tyr605)

at Human:Y605, Mouse:Y606

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source : Polyclonal, Rabbit, IgG

Dilution : IHC 1:50-200, WB 1:500-2000. IF 1:50-200

Purification: The antibody was affinity-purified from rabbit serum by affinity-chromatography

using specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 108kD

1/3



Background:

This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. This gene is expressed in some human cancer cell lines and has been implicated in carcinogenesis. [provided by RefSeq, Jul 2008],

Function:


catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,function:Receptor for members of the ephrin-A family. Binds with a low affinity to ephrin-A1.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. Ephrin receptor subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,similarity:Contains 2 fibronectin type-III domains.,tissue specificity:Overexpressed in several carcinomas.,

Subcellular Location:


Cell membrane; Single-pass type I membrane protein.

Expression : Overexpressed in several carcinomas.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



Western Blot analysis of 1 Hela treated with LPS, 2 Hela,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000