

## TPOR (Phospho Tyr626) rabbit pAb

|                              |   |
|------------------------------|---|
| <b>Catalog No :</b>          | YP1535  |
| <b>Reactivity :</b>          | Human;Rat;Mouse;  |
| <b>Applications :</b>        | WB  |
| <b>Target :</b>              | CD110   |
| <b>Fields :</b>              | >>Cytokine-cytokine receptor interaction;>>JAK-STAT signaling pathway                                     |
| <b>Gene Name :</b>           | MPL TPOR  |
| <b>Protein Name :</b>        | TPOR (Tyr626)   |
| <b>Human Gene Id :</b>       | 4352  |
| <b>Human Swiss Prot No :</b> | P40238  |
| <b>Mouse Gene Id :</b>       | 17480   |
| <b>Mouse Swiss Prot No :</b> | Q08351  |
| <b>Immunogen :</b>           | Synthesized phosho peptide around human TPOR (Tyr626)   |
| <b>Specificity :</b>         | This antibody detects endogenous levels of Human TPOR (phospho-Tyr626)                                    |
| <b>Formulation :</b>         | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                   |
| <b>Source :</b>              | Polyclonal, Rabbit,IgG  |
| <b>Dilution :</b>            | WB 1:1000-2000  |
| <b>Purification :</b>        | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| <b>Concentration :</b>       | 1 mg/ml   |

|                               |  |
|-------------------------------|--|
| <b>Storage Stability :</b>    | -15°C to -25°C/1 year(Do not lower than -25°C)   |
| <b>Observed Band :</b>        | 69,40kD  |
| <b>Cell Pathway :</b>         | Cytokine-cytokine receptor interaction;Jak_STAT;   |
| <b>Background :</b>           | <p>In 1990 an oncogene, v-mpl, was identified from the murine myeloproliferative leukemia virus that was capable of immortalizing bone marrow hematopoietic cells from different lineages. In 1992 the human homologue, named, c-mpl, was cloned. Sequence data revealed that c-mpl encoded a protein that was homologous with members of the hematopoietic receptor superfamily. Presence of anti-sense oligodeoxynucleotides of c-mpl inhibited megakaryocyte colony formation. The ligand for c-mpl, thrombopoietin, was cloned in 1994. Thrombopoietin was shown to be the major regulator of megakaryocytopoiesis and platelet formation. The protein encoded by the c-mpl gene, CD110, is a 635 amino acid transmembrane domain, with two extracellular cytokine receptor domains and two intracellular cytokine receptor box motifs . TPO-R deficient mice were severely thrombocytopenic, emphasizing the important</p>            |
| <b>Function :</b>             | <p>caution:It is uncertain whether Met-1 or Met-8 is the initiator.,disease:Defects in MPL are a cause of congenital amegakaryocytic thrombocytopenia (CAMT) [MIM:604498]. CAMT is a disease characterized by isolated thrombocytopenia and megakaryocytopenia with no physical anomalies.,domain:The box 1 motif is required for JAK interaction and/or activation.,domain:The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding.,function:Receptor for thrombopoietin. May represent a regulatory molecule specific for TPO-R-dependent immune responses.,similarity:Belongs to the type I cytokine receptor family. Type 1 subfamily.,similarity:Contains 2 fibronectin type-III domains.,subunit:Interacts with ATXN2L.,tissue specificity:Expressed at a low level in a large number of cells of hematopoietic origin. Isoform 1 and</p> |
| <b>Subcellular Location :</b> | Cell membrane ; Single-pass type I membrane protein. Golgi apparatus . Cell surface .  |
| <b>Expression :</b>           | Expressed at a low level in a large number of cells of hematopoietic origin. Isoform 1 and isoform 2 are always found to be coexpressed.   |

## Products Images