

TNK1 (Phospho Tyr277) rabbit pAb

| | |
|------------------------------|---|
| Catalog No : | YP1532 |
| Reactivity : | Human;Rat;Mouse; |
| Applications : | WB |
| Target : | TNK1 |
| Gene Name : | TNK1 |
| Protein Name : | TNK1 (Tyr277) |
| Human Gene Id : | 8711 |
| Human Swiss Prot No : | Q13470 |
| Mouse Gene Id : | 83813 |
| Mouse Swiss Prot No : | Q99ML2 |
| Immunogen : | Synthesized phospho peptide around human TNK1 (Tyr277) |
| Specificity : | This antibody detects endogenous levels of Human TNK1 (phospho-Tyr277) |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | WB 1:1000-2000 |
| Purification : | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |

Observed Band : 75kD

Background : The protein encoded by this gene belongs to the tyrosine protein kinase family. Tyrosine protein kinases are important regulators of intracellular signal transduction pathways, mediating cellular proliferation, survival, and development. This gene is highly expressed in fetal tissues and at lower levels in few adult tissues, thus may function in signaling pathways utilized broadly during fetal development, and more selectively in adult tissues. It plays a negative regulatory role in the Ras-Raf1-MAPK pathway, and knockout mice have been shown to develop spontaneous tumors, suggesting a role as a tumor suppressor gene. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2011],

Function : catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,function:Involved in negative regulation of cell growth. Has tumor suppressor properties. Plays a negative regulatory role in the Ras-MAPK pathway. May function in signaling pathways utilized broadly during fetal development and more selectively in adult tissues and in cells of the lymphohematopoietic system. Could specifically be involved in phospholipid signal transduction.,PTM:Autophosphorylated on tyrosine residues.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 SH3 domain.,subunit:Interacts with the SH3 domain of PLCG1 via its Pro-rich domain.,tissue specificity:Expressed in all umbilical cord blood, bone marrow and adult blood cell sub-populations and in several leukemia cell lines. Highly expr

Subcellular Location : Cytoplasm . Membrane ; Peripheral membrane protein .

Expression : Expressed in all umbilical cord blood, bone marrow and adult blood cell sub-populations and in several leukemia cell lines. Highly expressed in fetal blood, brain, lung, liver and kidney. Detected at lower levels in adult prostate, testis, ovary, small intestine and colon. Not expressed in adult lung, liver, kidney or brain.

Products Images