

Ku-70 (Acetyl Lys539) Polyclonal Antibody

Catalog No :	YK0032
Reactivity :	Human;Rat;Mouse;
Applications :	WB;IHC;IF;ELISA
Target :	Ku70/XRCC6
Fields :	>>Non-homologous end-joining
Gene Name :	XRCC6
Protein Name :	X-ray repair cross-complementing protein 6
Human Gene Id :	2547
Human Swiss Prot No :	P12956
Mouse Swiss Prot No :	P23475
Immunogen :	Synthesized acetyl-peptide derived from the human Ku-70 around the acetylation site of K539.
Specificity :	Acetyl-Ku-70 (K539) Polyclonal Antibody detects endogenous levels of Ku-70 protein only when acetylated at K539.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC: 1:100-300 ELISA: 1:20000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 70kD

Cell Pathway : Non-homologous end-joining;


Background : The p70/p80 autoantigen is a nuclear complex consisting of two subunits with molecular masses of approximately 70 and 80 kDa. The complex functions as a single-stranded DNA-dependent ATP-dependent helicase. The complex may be involved in the repair of nonhomologous DNA ends such as that required for double-strand break repair, transposition, and V(D)J recombination. High levels of autoantibodies to p70 and p80 have been found in some patients with systemic lupus erythematosus. [provided by RefSeq, Jul 2008],

Function : developmental stage:Expression does not increase during promyelocyte differentiation.,disease:Individuals with systemic lupus erythematosus (SLE) and related disorders produce extremely large amounts of autoantibodies to p70 and p86. Existence of a major autoantigenic epitope or epitopes on the C-terminal 190 amino acids of p70 containing the leucine repeat. The majority of autoantibodies to p70 in most sera from patients with SLE seem to be reactive with this region.,function:Single stranded DNA-dependent ATP-dependent helicase. Has a role in chromosome translocation. The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner. It works in the 3'-5' direction. Binding to DNA may be mediated by p70. Involved in DNA nonhomologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination. The Ku p70/p86


Subcellular Location : Nucleus . Chromosome .

Expression : Brain,Cervix carcinoma,Epithelium,Heart,Hepatocyte,Kidney,Liver,Lun


Products Images



Li, H., Ma, L., Bian, X. et al. FK228 sensitizes radioresistant small cell lung cancer cells to radiation. Clin Epigenet 13, 41 (2021).



Western Blot analysis of Jurkat cells using Acetyl-Ku-70 (K539) Polyclonal Antibody. Antibody was diluted at 1:500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-breast, antibody was diluted at 1:100