

**AR (Acetyl Lys633) Polyclonal Antibody**

|                              |   |
|------------------------------|---|
| <b>Catalog No :</b>          | YK0077  |
| <b>Reactivity :</b>          | Human:K633;Mouse:K613;Rat:K616  |
| <b>Applications :</b>        | WB;ELISA  |
| <b>Target :</b>              | Androgen Receptor   |
| <b>Fields :</b>              | >>Oocyte meiosis;>>Pathways in cancer;>>Chemical carcinogenesis - receptor activation;>>Prostate cancer               |
| <b>Gene Name :</b>           | AR DHTR NR3C4   |
| <b>Protein Name :</b>        | Androgen receptor (Dihydrotestosterone receptor) (Nuclear receptor subfamily 3 group C member 4)                      |
| <b>Human Gene Id :</b>       | 367   |
| <b>Human Swiss Prot No :</b> | P10275  |
| <b>Mouse Swiss Prot No :</b> | P19091  |
| <b>Rat Swiss Prot No :</b>   | P15207  |
| <b>Immunogen :</b>           | Synthetic Acetyl peptide from human protein at AA range: 633  |
| <b>Specificity :</b>         | This antibody detects endogenous levels of AR at Human:K633;Mouse:K613;Rat:K616, It doesn't react with total protein. |
| <b>Formulation :</b>         | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| <b>Source :</b>              | Polyclonal, Rabbit,IgG  |
| <b>Dilution :</b>            | WB 1:500-2000, ELISA 1:10000-20000  |
| <b>Purification :</b>        | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |

**Concentration :** 1 mg/ml

---

**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)

---

**Observed Band :** 100kD

---

**Cell Pathway :** Oocyte meiosis;Pathways in cancer;Prostate cancer;

---

**Background :** The androgen receptor gene is more than 90 kb long and codes for a protein that has 3 major functional domains: the N-terminal domain, DNA-binding domain, and androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its protein. Expansion of the polyglutamine tract from the normal 9-34 repeats to the pathogenic 38-62 repeats causes spinal bulbar muscular atrophy (Kennedy disease). Mutations in this gene are also associated with complete androgen insensitivity (CAIS). Two alternatively spliced variants encoding distinct isoform

---

**Function :** disease:Defects in AR are the cause of androgen insensitivity syndrome (AIS) [MIM:300068]; previously known as testicular feminization syndrome (TFM). AIS is an X-linked recessive form of pseudohermaphroditism due end-organ resistance to androgen. Affected males have female external genitalia, female breast development, blind vagina, absent uterus and female adnexa, and abdominal or inguinal testes, despite a normal 46,XY karyotype.,disease:Defects in AR are the cause of androgen insensitivity syndrome partial (PAIS) [MIM:312300]; also known as Reifenstein syndrome. PAIS is characterized by hypospadias, hypogonadism, gynecomastia, genital ambiguity, normal XY karyotype, and a pedigree pattern consistent with X-linked recessive inheritance. Some patients present azoospermia or severe oligospermia without other clinical manifestations.,disease:Defects in AR are the cause of spinal and bulb

---


**Subcellular Location :** Nucleus . Cytoplasm . Detected at the promoter of target genes (PubMed:25091737). Predominantly cytoplasmic in unligated form but translocates to the nucleus upon ligand-binding. Can also translocate to the nucleus in unligated form in the presence of RACK1 . .

---

**Expression :** [Isoform 2]: Mainly expressed in heart and skeletal muscle. ; [Isoform 3]: Expressed in basal and stromal cells of the prostate (at protein level).

---

## Products Images



Western blot analysis of mouse-lung lysate, antibody was diluted at 500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000