

HMGN1/2/3/4 (Acetyl Lys27/K33/K31) rabbit pAb

Catalog No :	YK0146
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	HMGN1/2/3/4
Gene Name :	HMGN1 HMG14
Protein Name :	HMGN1/2/3/4 (Acetyl Lys27/K33/K31)
Human Gene Id :	3150
Human Swiss Prot No :	P05114/P05204/Q15651/O00479
Mouse Gene Id :	100044391
Mouse Swiss Prot No :	P18608
Immunogen :	Synthesized peptide derived from human HMGN1/2/3/4 (Acetyl Lys27/K33/K31)
Specificity :	This antibody detects endogenous levels of Human,Mouse,Rat HMGN1/2/3/4 (Acetyl Lys27/K33/K31)
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:1000-2000 ELISA 1:5000-20000
Purification :	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 12kD

Background : The protein encoded by this gene binds nucleosomal DNA and is associated with transcriptionally active chromatin. Along with a similar protein, HMG17, the encoded protein may help maintain an open chromatin configuration around transcribable genes. [provided by RefSeq, Aug 2011],

Function : function: Binds to the inner side of the nucleosomal DNA thus altering the interaction between the DNA and the histone octamer. May be involved in the process which maintains transcribable genes in a unique chromatin conformation. Inhibits the phosphorylation of nucleosomal histones H3 and H2A by RPS6KA5/MSK1 and RPS6KA3/RSK2., mass spectrometry: PubMed:10739259, PTM: Phosphorylation on Ser-21 and Ser-25 weakens binding to nucleosomes and increases the rate of H3 phosphorylation (By similarity). Phosphorylation favors cytoplasmic localization., RNA editing: Partially edited. A new initiator methionine may be created by a single uridine insertion in the 5'-UTR, causing an N-terminal extension of 45 amino acids. The existence of the RNA edited version is supported by direct protein sequencing by MS/MS of the following peptides specific to that version: 23-31 and 40-48. The RNA edited version is

Subcellular Location : Nucleus. Cytoplasm. Cytoplasmic enrichment upon phosphorylation. The RNA edited version localizes to the nucleus.

Products Images