

CD131 (PN0305) Nb-FC recombinant antibody

Catalog No :	YA0066
Reactivity :	Human
Applications :	ELISA
Target :	_CD131
Gene Name :	CSF2RB IL3RB IL5RB
Protein Name :	Cytokine receptor common subunit beta (CDw131) (GM-CSF/IL-3/IL-5 receptor common beta subunit) (CD antigen CD131)
Human Gene Id :	1439
Human Swiss Prot No :	P32927
Immunogen :	Purified recombinant Human CD131
Specificity :	This recombinant monoclonal antibody can detects endogenous levels of CD131 protein.
Formulation :	Phosphate-buffered solution
Source :	Camel, chimeric fusion of Nanobody (VHH) and mouse $\mbox{IgG1}$ Fc domain , recombinantly produced from 293F cell
Dilution :	ELISA 1:5000-100000
Purification :	Recombinant Expression and Affinity purified
Concentration :	Please check the information on the tube
Storage Stability :	-15°C to -25°C/1 year(Avoid freeze / thaw cycles)
Background :	CD131, also known as the IL-3R common β subunit , is a 95-120 kD type I transmembrane glycoprotein and belongs to the Ig superfamily. The common β subunit associates with the specific g subunits of IL-3 receptor. IL-5 receptor and



GM-CSF receptor to form high affinity receptors for these cytokines. These cytokine receptors are expressed by neutrophils, eosinophils, monocytes, endothelial cells, fibroblasts and hematopoietic progenitor cells and play a crucial role in growth/activation of eosinophils and in the inflammatory response. The 1C1 antibody is a non-blocking antibody.

Function:

Cell surface receptor that plays a role in immune response and controls the production and differentiation of hematopoietic progenitor cells into lineagerestricted cells. Acts by forming an heterodimeric receptor through interaction with different partners such as IL3RA, IL5RA or CSF2RA (PubMed:1495999). In turn, participates in various signaling pathways including interleukin-3, interleukin-5 and granulocyte-macrophage colony-stimulating factor/CSF2 pathways. In unstimulated conditions, interacts constitutively with JAK1 and ligand binding leads to JAK1 stimulation and subsequent activation of the JAK-STAT pathway (PubMed:9516124).

Subcellular Location :

Membrane; Single-pass type I membrane protein.

Products Images

