

CD23 (PN0254) Nb-FC recombinant antibody

YA0198 Catalog No:

Reactivity: Human

ELISA;FCM **Applications:**

Target: CD23

Gene Name: FCER2 CD23A CLEC4J FCE2 IGEBF

Protein Name: Low affinity immunoglobulin epsilon Fc receptor (BLAST-2) (C-type lectin

domain family 4 member J) (Fc-epsilon-RII) (Immunoglobulin E-binding factor)

(Lymphocyte IgE receptor) (CD antigen CD23) [Cleav

Human Gene Id: 2208

Human Swiss Prot

No:

Purified recombinant Human CD23 Immunogen:

P06734

This recombinant monoclonal antibody can detects endogenous levels of CD23 **Specificity:**

protein.

Formulation: Phosphate-buffered solution

Source: Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain,

recombinantly produced from 293F cell

Dilution: ELISA 1:5000-100000 FCM 1-2µg/Test

Purification: Recombinant Expression and Affinity purified

Concentration: Please check the information on the tube

-15°C to -25°C/1 year(Avoid freeze / thaw cycles) **Storage Stability:**

Cell Pathway: Hematopoietic cell lineage;

1/2



Background:

The protein encoded by This gene is a B-cell specific antigen, and a low-affinity receptor for IgE. It has essential roles in B cell growth and differentiation, and the regulation of IgE production. This protein also exists as a soluble secreted form, then functioning as a potent mitogenic growth factor. Alternatively spliced transcript variants encoding different isoforms have been described for This gene. [provided by Ref Seq, Jul 2011]

Function:

This receptor has essential roles in the regulation of IgE production and in the differentiation of B-cells (it is a B-cell-specific antigen).,miscellaneous:There are two kinds of Fc receptors for IgE, which differ in both structure and high affinity receptors on basophils and mast cells and low affinity receptors on lymphocytes and monocytes.,online information:CD23,PTM:N- and O-glycosylated.,similarity:Contains 1 C-type lectin domain.,subcellular location:Also exists as a soluble excreted form.,

Subcellular Location:

Cell membrane; Single-pass type II membrane protein. Cell membrane; Lipidanchor. Secreted. Also exists as a soluble excreted form, sCD23.

Expression : B-cell,Blood

Products Images

