

CD269/BCMA (PN0553) Nb-FC recombinant antibody

| Catalog No : | YA0222 |
|--------------------------|---|
| Reactivity : | Human |
| Applications : | FCM;ELISA |
| Target : | CD269/BCMA |
| Gene Name : | TNFRSF17 BCM BCMA |
| Protein Name : | Tumor necrosis factor receptor superfamily member 17 (B-cell maturation protein) (CD antigen CD269) |
| Human Gene Id : | 608 |
| Human Swiss Prot No : | Q02223 |
| Immunogen : | Purified recombinant Human CD269 |
| Specificity : | This recombinant monoclonal antibody can detects endogenous levels of CD269/BCMA protein. |
| Formulation : | Phosphate-buffered solution |
| Source : | Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell |
| Dilution : | ELISA 1:5000-100000;FCM 1-2µg/Test |
| Purification : | Recombinant Expression and Affinity purified |
| Concentration : | Please check the information on the tube |
| Storage Stability : | -15°C to -25°C/1 year(Avoid freeze / thaw cycles) |
| Cell Pathway : | Cytokine-cytokine receptor interaction;Intestinal immune network for IgA production; |



| Background : | The protein encoded byThis gene is a member of the TNF-receptor superfamily.This receptor is preferentially expressed in mature B lymphocytes, and may be important for B cell development and autoimmune response.This receptor has been shown to specifically bind to the tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B/TALL-1/BAFF), and to lead to NF-kappaB and MAPK8/JNK activation.This receptor also binds to various TRAF family members, and thus may transduce signals for cell survival and proliferation. [provided by RefSeq, Jul 2008] |
|---------------------------|--|
| Function : | disease:A chromosomal aberration involving TNFRSF17 is found in a form of T- cell acute lymphoblastic leukemia (T-ALL). Translocation t(4;16)(q26;p13) with IL2.,Receptor for TNFSF13B/BLyS/BAFF and TNFSF13/APRIL. Promotes B-cell survival and plays a role in the regulation of humoral immunity. Activates NF- kappa-B and JNK.,similarity:Contains 1 TNFR-Cys repeat.,subcellular location:Perinuclear Golgi-like structures.,subunit:Associates with TRAF1, TRAF2, TRAF3, TRAF5 and TRAF6.,tissue specificity:Expressed in mature B- cells, but not in T-cells or monocytes., |
| Subcellular Location : | Cell membrane; Single-pass type III membrane protein. Endomembrane system; Single-pass type III membrane protein. Perinuclear Golgi-like structures. |
| Expression : | Expressed in mature B-cells, but not in T-cells or monocytes. |

Products Images