

CD9 (PN0189) Nb-FC recombinant antibody

Catalog No: YA0554

Reactivity: Human

Applications: ELISA;FCM

Target: CD9

Gene Name: CD9 MIC3 TSPAN29 GIG2

Protein Name: CD9 antigen (5H9 antigen) (Cell growth-inhibiting gene 2 protein) (Leukocyte

antigen MIC3) (Motility-related protein) (MRP-1) (Tetraspanin-29) (Tspan-29)

(p24) (CD antigen CD9)

Human Gene Id: 928

Human Swiss Prot

No:

Immunogen: Purified recombinant Human CD9

P21926

Specificity: This recombinant monoclonal antibody can detects endogenous levels of CD9

protein.

Formulation : Phosphate-buffered solution

Source: Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain,

recombinantly produced from 293F cell

Dilution: ELISA 1:5000-100000 FCM 1-2µg/Test

Purification: Recombinant Expression and Affinity purified

Concentration: Please check the information on the tube

Storage Stability: -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

Cell Pathway: Hematopoietic cell lineage;

1/2

Background:

This gene encodes a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Tetraspanins are cell surface glycoproteins with four transmembrane domains that form multimeric complexes with other cell surface proteins. The encoded protein functions in many cellular processes including differentiation, adhesion, and signal transduction, and expression of This gene plays a critical role in the suppression of cancer cell motility and metastasis. [provided by RefSeq, Jan 2011]

Function:

Involved in platelet activation and aggregation. Regulates paranodal junction formation. Involved in cell adhesion, cell motility and tumor metastasis. Required for sperm-egg fusion.,PTM:Protein exists in three forms with molecular masses between 22 and 27 kDa, and is known to carry covalently linked fatty acids.,similarity:Belongs to the tetraspanin (TM4SF) family.,subunit:Forms both disulfide-linked homodimers and higher homooligomers as well as heterooligomers with other members of the tetraspanin family. Associates with CR2/CD21 and with PTGFRN/CD9P1. Interacts directly with IGSF8.,tissue specificity:Expressed by a variety of hematopoietic and epithelial cells.,

Subcellular Location:

Cell membrane ; Multi-pass membrane protein . Membrane ; Multi-pass membrane protein . Secreted, extracellular exosome . Present at the cell surface of oocytes. Accumulates in the adhesion area between the sperm and egg following interaction between IZUMO1 and its receptor IZUMO1R/JUNO. .

Expression:

Detected in platelets (at protein level) (PubMed:1964571). Expressed by a variety of hematopoietic and epithelial cells (PubMed:1964571).

Products Images

