

## CD63 (PN0219) Nb-FC recombinant antibody

| Catalog No :             | YA0436  |
|--------------------------|---|
| Reactivity :             | Human   |
| Applications :           | ELISA   |
| Target :                 | CD63  |
| Gene Name :              | CD63 MLA1 TSPAN30   |
| Protein Name :           | CD63 antigen (Granulophysin) (Lysosomal-associated membrane protein 3) (LAMP-3) (Lysosome integral membrane protein 1) (Limp1) (Melanoma-associated antigen ME491) (OMA81H) (Ocular melanoma-associated |
| Human Gene Id :          | 967   |
| Human Swiss Prot<br>No : | P08962  |
| Immunogen :              | Purified recombinant Human CD63   |
| Specificity :            | This recombinant monoclonal antibody can detects endogenous levels of CD63 protein.   |
| Formulation :            | Phosphate-buffered solution   |
| Source :                 | Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell   |
| Dilution :               | ELISA 1:5000-100000   |
| Purification :           | Recombinant Expression and Affinity purified  |
| Concentration :          | Please check the information on the tube  |
| Storage Stability :      | -15°C to -25°C/1 year(Avoid freeze / thaw cycles)   |
| Cell Pathway :           | Lysosome;   |



| Background :              | The protein encoded byThis gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. The encoded protein is a cell surface glycoprotein that is known to complex with integrins. It may function as a blood platelet activation marker. Deficiency of This protein is associated with Hermansky-Pudlak syndrome. AlsoThis gene has been associated with tumor progression. Alternative splicing results in multiple transcript variants encoding different protein isoforms. [provided by RefSeq, Apr 2012]   |
|---------------------------|---|
| Function :                | This antigen is associated with early stages of melanoma tumor progression.<br>May play a role in growth regulation.,miscellaneous:Lack of expression of CD63 in<br>platelets has been observed in a patient with Hermansky-Pudlak syndrome<br>(HPS). Hermansky-Pudlak syndrome (HPS) is a genetically heterogeneous, rare,<br>autosomal recessive disorder characterized by oculocutaneous albinism,<br>bleeding due to platelet storage pool deficiency, and lysosomal storage defects.<br>This syndrome results from defects of diverse cytoplasmic organelles including<br>melanosomes, platelet dense granules and lysosomes. Ceroid storage in the<br>lungs is associated with pulmonary fibrosis, a common cause of premature death<br>in individuals with HPS.,similarity:Belongs to the tetraspanin (TM4SF)<br>family.,subcellular location:Also found in Weibel-Palade bodies of endothelial<br>cells. Located in platelet dense granules.,tissue specifici |
| Subcellular<br>Location : | Cell membrane ; Multi-pass membrane protein . Lysosome membrane ; Multi-<br>pass membrane protein . Late endosome membrane ; Multi-pass membrane<br>protein . Endosome, multivesicular body . Melanosome . Secreted, extracellular<br>exosome . Cell surface . Also found in Weibel-Palade bodies of endothelial cells<br>(PubMed:10793155). Located in platelet dense granules (PubMed:7682577).<br>Detected in a subset of pre-melanosomes. Detected on intralumenal vesicles<br>(ILVs) within multivesicular bodies (PubMed:21962903)  |
| Expression :              | Detected in platelets (at protein level). Dysplastic nevi, radial growth phase primary melanomas, hematopoietic cells, tissue macrophages.  |

## Products Images



