

## CD80 Monoclonal Antibody

<b>Catalog No :</b>	YM0136
<b>Reactivity :</b>	Human
<b>Applications :</b>	IHC;IF;ELISA
<b>Target :</b>	CD80
<b>Fields :</b>	>>Cell adhesion molecules;>>Toll-like receptor signaling pathway;>>Intestinal immune network for IgA production;>>Type I diabetes mellitus;>>Autoimmune thyroid disease;>>Systemic lupus erythematosus;>>Rheumatoid arthritis;>>Allograft rejection;>>Graft-versus-host disease;>>Viral myocarditis
<b>Gene Name :</b>	CD80
<b>Protein Name :</b>	T-lymphocyte activation antigen CD80
<b>Human Gene Id :</b>	941
<b>Human Swiss Prot No :</b>	P33681
<b>Mouse Swiss Prot No :</b>	Q00609
<b>Immunogen :</b>	Purified recombinant fragment of CD80 expressed in E. Coli.
<b>Specificity :</b>	CD80 Monoclonal Antibody detects endogenous levels of CD80 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Monoclonal, Mouse
<b>Dilution :</b>	IHC 1:200 - 1:1000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.
<b>Purification :</b>	Affinity purification
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

**Cell Pathway :** Cell adhesion molecules (CAMs);Toll\_Like;Intestinal immune network for IgA production;Type I diabetes mellitus;Autoimmune thyroid disease;Systemic lupus erythematosus;Allograft rejection;Graft-versus-

**P References :** 1. Transplant Proc. 2008 Oct;40(8):2729-33.  
2. Nat Med. 2007 Dec;13(12):1440-9.


**Background :** The protein encoded by this gene is a membrane receptor that is activated by the binding of CD28 or CTLA-4. The activated protein induces T-cell proliferation and cytokine production. This protein can act as a receptor for adenovirus subgroup B and may play a role in lupus neuropathy. [provided by RefSeq, Aug 2011],

**Function :** function:Involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is induced by the binding of CD28 or CTLA-4 to this receptor.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,subunit:Interacts with adenovirus subgroup B fiber proteins and acts as a receptor for these viruses.,tissue specificity:Expressed on activated B-cells, macrophages and dendritic cells.,


**Subcellular Location :** Membrane; Single-pass type I membrane protein.

**Expression :** Expressed on activated B-cells, macrophages and dendritic cells.

## Products Images



Immunohistochemistry analysis of paraffin-embedded human Tonsil tissues with AEC staining using CD80 Monoclonal Antibody.



Confocal immunofluorescence analysis of BCBL-1 cells using CD80 Monoclonal Antibody (green), showing membrane localization. Blue: DRAQ5 fluorescent DNA dye.