

## Crk II Monoclonal Antibody

<b>Catalog No :</b>	YM0167
<b>Reactivity :</b>	Human
<b>Applications :</b>	WB;IHC;IF;FCM;ELISA
<b>Target :</b>	Crk II
<b>Fields :</b>	>>MAPK signaling pathway;>>ErbB signaling pathway;>>Rap1 signaling pathway;>>Chemokine signaling pathway;>>Focal adhesion;>>Fc gamma R-mediated phagocytosis;>>Neurotrophin signaling pathway;>>Regulation of actin cytoskeleton;>>Insulin signaling pathway;>>Growth hormone synthesis, secretion and action;>>Bacterial invasion of epithelial cells;>>Shigellosis;>>Yersinia infection;>>Human cytomegalovirus infection;>>Human immunodeficiency virus 1 infection;>>Pathways in cancer;>>MicroRNAs in cancer;>>Renal cell carcinoma;>>Chronic myeloid leukemia
<b>Gene Name :</b>	CRK
<b>Protein Name :</b>	Adapter molecule crk
<b>Human Gene Id :</b>	1398
<b>Human Swiss Prot No :</b>	P46108
<b>Mouse Swiss Prot No :</b>	Q64010
<b>Immunogen :</b>	Purified recombinant fragment of human Crk II expressed in E. Coli.
<b>Specificity :</b>	Crk II Monoclonal Antibody detects endogenous levels of Crk II protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Monoclonal, Mouse
<b>Dilution :</b>	WB 1:500 - 1:2000. IHC 1:200 - 1:1000. IF 1:200 - 1:1000. Flow cytometry: 1:200 - 1:400. ELISA: 1:10000. Not yet tested in other applications.
	Affinity purification

**Storage Stability :** -15°C to -25°C/1 year (Do not lower than -25°C)

**Molecular weight :** 34kD

**Cell Pathway :** MAPK\_ERK\_Growth;MAPK\_G\_Protein;ErbB\_HER;Chemokine;Focal adhesion;Fc gamma R-mediated phagocytosis;Neurotrophin;Regulates Actin and Cytoskeleton;Insulin\_Receptor;Pathways in cancer;Renal cell carcinoma

**P References :**

1. Seikagaku. 2009 May;81(5):361-76.
2. Mol Cancer Res. 2009 Sep;7(9):1582-92.


**Background :** This gene encodes a member of an adapter protein family that binds to several tyrosine-phosphorylated proteins. The product of this gene has several SH2 and SH3 domains (src-homology domains) and is involved in several signaling pathways, recruiting cytoplasmic proteins in the vicinity of tyrosine kinase through SH2-phosphotyrosine interaction. The N-terminal SH2 domain of this protein functions as a positive regulator of transformation whereas the C-terminal SH3 domain functions as a negative regulator of transformation. Two alternative transcripts encoding different isoforms with distinct biological activity have been described. [provided by RefSeq, Jul 2008],

**Function :** domain:The C-terminal SH3 domain function as a negative modulator for transformation and the N-terminal SH3 domain appears to function as a positive regulator for transformation.,domain:The SH2 domain mediates interaction with SHB.,function:The Crk-I and Crk-II forms differ in their biological activities. Crk-II has less transforming activity than Crk-I. Crk-II mediates attachment-induced MAPK8 activation, membrane ruffling and cell motility in a Rac-dependent manner. Involved in phagocytosis of apoptotic cells and cell motility via its interaction with DOCK1 and DOCK4.,PTM:Phosphorylated on Tyr-221 upon cell adhesion. Results in the negative regulation of the association with SH2- and SH3-binding partners, possibly by the formation of an intramolecular interaction of phosphorylated Tyr-221 with the SH2 domain. This leads finally to the down-regulation of the Crk signaling pathway.,PTM:P


**Subcellular Location :** Cytoplasm . Cell membrane . Translocated to the plasma membrane upon cell adhesion. .

**Expression :** Embryonic lung,Epithelium,Eye,Lung,Placenta,


## Products Images




Western Blot analysis using Crk II Monoclonal Antibody against HEK293 (1) and CRK-hlgGfc transfected HEK293 (2) cell lysate.



Immunohistochemistry analysis of paraffin-embedded intima cancer tissues with DAB staining using Crk II Monoclonal Antibody.



Immunofluorescence analysis of 3T3-L1 cells using Crk II Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.



Flow cytometric analysis of Hela cells using Crk II Monoclonal Antibody (blue) and negative control (red).

