


HDAC4 Monoclonal Antibody

Catalog No :	YM1041
Reactivity :	Human;Mouse;Rat
Applications :	WB
Target :	HDAC4
Fields :	>>Apelin signaling pathway;>>Neutrophil extracellular trap formation;>>Alcoholism;>>Viral carcinogenesis;>>MicroRNAs in cancer
Gene Name :	HDAC4
Protein Name :	Histone deacetylase 4
Human Gene Id :	9759
Human Swiss Prot No :	P56524
Mouse Gene Id :	208727
Mouse Swiss Prot No :	Q6NZM9
Rat Gene Id :	363287
Rat Swiss Prot No :	Q99P99
Immunogen :	Purified recombinant human HDAC4 (N-terminus) protein fragments expressed in E.coli.
Specificity :	HDAC4 Monoclonal Antibody detects endogenous levels of HDAC4 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	WB 1:1000 - 1:2000. Not yet tested in other applications.

Purification :	Affinity purification
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	119kD
Cell Pathway :	Protein_Acetylation
Background :	<p>Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a multiprotein complex with RbAp48 and HDAC3. [provided by RefSeq, Jul 2008],</p>
Function :	<p>catalytic activity:Hydrolysis of an N(6)-acetyl-lysine residue of a histone to yield a deacetylated histone.,domain:The nuclear export sequence mediates the shuttling between the nucleus and the cytoplasm.,function:Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Involved in muscle maturation via its interaction with the myocyte enhancer factors such as MEF2A, MEF2C and MEF2D.,PTM:Phosphorylated by CaMK4 at Ser-246, Ser-467 and Ser-632. Phosphorylation at other residues is required for the interaction with 14-3-3.,PTM:Sumoylation on Lys-559 is promoted by the E3 SUMO-protein lig</p>
Subcellular Location :	<p>Nucleus. Cytoplasm. Shuttles between the nucleus and the cytoplasm. Upon muscle cells differentiation, it accumulates in the nuclei of myotubes, suggesting a positive role of nuclear HDAC4 in muscle differentiation. The export to cytoplasm depends on the interaction with a 14-3-3 chaperone protein and is due to its phosphorylation at Ser-246, Ser-467 and Ser-632 by CaMK4 and SIK1. The nuclear localization probably depends on sumoylation. Interaction with SIK3 leads to HDAC4 retention in the cytoplasm (By similarity). .</p>
Expression :	Ubiquitous.

Products Images



Western Blot analysis using HDAC4 Monoclonal Antibody against Jurkat, K562 cell lysate.