

PRMT4/CARM1 mouse mAb

Catalog No :	YM1215
Reactivity :	Human;Mouse
Applications :	WB;IP
Target :	CARM1
Fields :	>>Endocrine resistance
Gene Name :	carm1
Human Gene Id :	10498
Human Swiss Prot No :	Q86X55
Mouse Swiss Prot No :	Q9WVG6
Immunogen :	Purified recombinant human PRMT4/CARM1 protein fragments expressed in E.coli.
Specificity :	This antibody detects endogenous levels of PRMT4/CARM1 and does not cross-react with related proteins.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	wb 1:200-1:500
Purification :	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	63kD


Background : This gene belongs to the protein arginine methyltransferase (PRMT) family. The encoded enzyme catalyzes the methylation of guanidino nitrogens of arginyl residues of proteins. The enzyme acts specifically on histones and other chromatin-associated proteins and is involved in regulation of gene expression. The enzyme may act in association with other proteins or within multi-protein complexes and may play a role in cell type-specific functions and cell lineage specification. A related pseudogene is located on chromosome 9. [provided by RefSeq, Aug 2013],

Function : catalytic activity:S-adenosyl-L-methionine + histone-arginine = S-adenosyl-L-homocysteine + histone-N(omega)-methyl-arginine.,function:Methylates (mono- and asymmetric dimethylation) the guanidino nitrogens of arginyl residues in several proteins involved in DNA packaging, transcription regulation, and mRNA stability. Recruited to promoters upon gene activation together with histone acetyltransferases from EP300/P300 and p160 families, methylates histone H3 at 'Arg-17' and activates transcription via chromatin remodeling. During nuclear hormone receptor activation and TCF7L2/TCF4 activation, acts synergically with EP300/P300 and either one of the p160 histone acetyltransferases NCOA1/SRC1, NCOA2/GRIP1 and NCOA3/ACTR or CTNNB1/beta-catenin to activate transcription. During myogenic transcriptional activation, acts together with NCOA3/ACTR as a coactivator for MEF2C. During monocyte inflam

Subcellular Location : Nucleus . Cytoplasm . Mainly nuclear during the G1, S and G2 phases of the cell cycle (PubMed:19843527). Cytoplasmic during mitosis, after breakup of the nuclear membrane (PubMed:19843527)..

Expression : Overexpressed in prostate adenocarcinomas and high-grade prostatic intraepithelial neoplasia.

Products Images



Western blot detection of PRMT4/CARM1 in HeLa,A431 and K562 cell lysates using PRMT4/CARM1 mouse mAb (1:200-1:500 diluted).Predicted band size:63KDa.Observed band size:63KDa.

Immunoprecipitation analysis of HeLa cell lysates using PRMT4/CARM1 mouse mAb.

