

KAP1/TIF1 β mouse mAb

Catalog No :	YM1235
Reactivity :	Human
Applications :	WB;ICC;IHC;IP
Target :	TIF1 β
Gene Name :	trim28
Human Gene Id :	10155
Human Swiss Prot No :	Q13263
Mouse Swiss Prot No :	Q62318
Immunogen :	Purified recombinant human KAP1 / TIF1 beta protein fragments expressed in E.coli.
Specificity :	This antibody detects endogenous levels of KAP1 / TIF1 beta and does not cross-react with related proteins.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	wb 1:1000 icc 1:100
Purification :	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	110kD
Background :	The protein encoded by this gene mediates transcriptional control by interaction

with the Kruppel-associated box repression domain found in many transcription factors. The protein localizes to the nucleus and is thought to associate with specific chromatin regions. The protein is a member of the tripartite motif family. This tripartite motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. [provided by RefSeq, Jul 2008],

Function :

domain:Contains one Pro-Xaa-Val-Xaa-Leu (PxVxL) motif, which is required for interaction with chromoshadow domains. This motif requires additional residues -7, -6, +4 and +5 of the central Val which contact the chromoshadow domain.,domain:The HP1 box is both necessary and sufficient for HP1 binding. The RING finger domain and the B-box domains mediate interaction with CEBPB. The PHD domain enhances the CEBPB transcriptional activity.,function:Forms a complex with a KRAB-domain transcription factor and increases the efficiency of KRAB-mediated repression. Silences transcription through an interaction with HP1 proteins. Acts as a corepressor of transcription for the KRAB zinc finger proteins and as a moderator of the repression activity. May play a role as a coactivator for CEBPB and NR3C1 in the transcriptional activation of the Alpha-1-acid glycoprotein gene.,PTM:Phosphorylated upon DNA


Subcellular Location :

Nucleus . Associated with centromeric heterochromatin during cell differentiation through CBX1. .


Expression :

Expressed in all tissues tested including spleen, thymus, prostate, testis, ovary, small intestine, colon and peripheral blood leukocytes.


Products Images




Western blot detection of KAP1 / TIF1 beta in 293T and HepG2 cell lysates using KAP1 / TIF1 beta mouse mAb (1:1000 diluted). Observed band size: 110KDa.




Immunocytochemistry staining of HeLa cells fixed with 4% Paraformaldehyde and using anti-KAP1 / TIF1 beta mouse mAb (dilution 1:100).



Immunoprecipitation analysis of Hela cell lysates using KAP1 / TIF1 beta mouse mAb.



IHC of paraffin-embedded human Spleen using anti-KAP1 / TIF1 beta diluted 1/500-1/1000.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).