

JMJD6(N-term) mouse mAb

Catalog No :	YM1204
Reactivity :	Human
Applications :	WB;ICC
Target :	PSR
Gene Name :	jmjd6
Human Gene Id :	23210
Human Swiss Prot No :	Q6NYC1
Mouse Swiss Prot No :	Q9ERI5
Immunogen :	Purified recombinant human JMJD6(N-terminus) fragments expressed in E.coli.
Specificity :	This antibody detects endogenous levels of JMJD6(N-terminus) and does not cross-react with related proteins.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	wb 1:1000 icc 1:200
Purification :	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	62kD
Background :	This gene encodes a nuclear protein with a JmjC domain. JmjC domain-containing proteins are predicted to function as protein hydroxylases or histone

demethylases. This protein was first identified as a putative phosphatidylserine receptor involved in phagocytosis of apoptotic cells; however, subsequent studies have indicated that it does not directly function in the clearance of apoptotic cells, and questioned whether it is a true phosphatidylserine receptor. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],

Function :

caution:Was initially thought to constitute the phosphatidylserine receptor, a receptor that mediates recognition of phosphatidylserine, a specific marker only present at the surface of apoptotic cells. Phosphatidylserine receptor probably participates in apoptotic cell phagocytosis. This protein was identified using phage display expressing mAb 217, an antibody that specifically recognizes phosphatidylserine receptor. However, its nuclear localization and the fact that mAb 217 antibody still recognizes the phosphatidylserine receptor in mice lacking JMJD6, strongly suggest that it does not constitute the receptor for phosphatidylserine and is not involved in apoptotic cell removal.,domain:The nuclear localization signal motifs are necessary and sufficient to target it into the nucleus.,function:Arginine demethylase which demethylates histone H3 at 'Arg-2' (H3R2me) and histone H4 at 'Arg'


Subcellular Location :

Nucleus, nucleoplasm . Nucleus, nucleolus . Cytoplasm . Mainly found throughout the nucleoplasm outside of regions containing heterochromatic DNA, with some localization in nucleolus. During mitosis, excluded from the nucleus and reappears in the telophase of the cell cycle. .


Expression :

Highly expressed in the heart, skeletal muscle and kidney. Expressed at moderate or low level in brain, placenta, lung, liver, pancreas, spleen, thymus, prostate, testis and ovary. Up-regulated in many patients with chronic pancreatitis. Expressed in nursing thymic epithelial cells.

Products Images



Western blot detection of JMJD6(N-terminus) in HeLa and HL-60 lysates using JMJD6(N-terminus) mouse mAb (1:1000 diluted). Predicted band size: 46KDa. Observed band size: 62KDa.



Immunocytochemistry of HeLa cells using anti-JMJD6(N-terminus) mouse mAb diluted 1:200.