


## PP1C mouse mAb

<b>Catalog No :</b>	YM1267
<b>Reactivity :</b>	Human
<b>Applications :</b>	WB
<b>Target :</b>	PP1C
<b>Fields :</b>	>>mRNA surveillance pathway;>>cGMP-PKG signaling pathway;>>cAMP signaling pathway;>>Oocyte meiosis;>>Cellular senescence;>>Adrenergic signaling in cardiomyocytes;>>Vascular smooth muscle contraction;>>Hippo signaling pathway;>>Focal adhesion;>>Platelet activation;>>Long-term potentiation;>>Dopaminergic synapse;>>Inflammatory mediator regulation of TRP channels;>>Regulation of actin cytoskeleton;>>Insulin signaling pathway;>>Oxytocin signaling pathway;>>Insulin resistance;>>Amphetamine addiction;>>Alcoholism;>>Herpes simplex virus 1 infection;>>Proteoglycans in cancer;>>Diabetic cardiomyopathy
<b>Gene Name :</b>	ppp1cc
<b>Human Gene Id :</b>	5501
<b>Human Swiss Prot No :</b>	P36873
<b>Mouse Swiss Prot No :</b>	P63087
<b>Immunogen :</b>	Purified recombinant human PP1C protein fragments expressed in E.coli.
<b>Specificity :</b>	This antibody detects endogenous levels of PP1C and does not cross-react with related proteins.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Monoclonal, Mouse
<b>Dilution :</b>	wb 1:500
<b>Purification :</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.

<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	38kD
<b>Cell Pathway :</b>	Oocyte meiosis;Vascular smooth muscle contraction;Focal adhesion;Long-term potentiation;Regulates Actin and Cytoskeleton;Insulin_Receptor;
<b>Background :</b>	<p>The protein encoded by this gene belongs to the protein phosphatase family, PP1 subfamily. PP1 is an ubiquitous serine/threonine phosphatase that regulates many cellular processes, including cell division. It is expressed in mammalian cells as three closely related isoforms, alpha, beta/delta and gamma, which have distinct localization patterns. This gene encodes the gamma isozyme. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2011],</p>
<b>Function :</b>	<p>catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate.,cofactor:Binds 1 iron ion per subunit.,cofactor:Binds 1 manganese ion per subunit.,enzyme regulation:The phosphatase activity of the PPP1R15A-PP1 complex toward EIF2S1 is specifically inhibited by Salubrinal, a drug that protects cells from endoplasmic reticulum stress.,function:Protein phosphatase 1 (PP1) is essential for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Involved in regulation of ionic conductances and long-term synaptic plasticity. May play an important role in dephosphorylating substrates such as the postsynaptic density-associated Ca(2+)/calmodulin dependent protein kinase II.,miscellaneous:Microcystin toxin is bound to Cys-273 through a thioether bond.,online information:The things we forget -Issue 32 of March 2003,similarity:Belon</p>
<b>Subcellular Location :</b>	<p>Cytoplasm . Nucleus. Nucleus, nucleolus . Nucleus, nucleoplasm . Nucleus speckle . Chromosome, centromere, kinetochore . Cleavage furrow . Midbody . Mitochondrion . Cytoplasm, cytoskeleton, microtubule organizing center . Colocalizes with SPZ1 in the nucleus (By similarity). Colocalizes with URI1 at mitochondrion (PubMed:17936702). Rapidly exchanges between the nucleolar, nucleoplasmic and cytoplasmic compartments (PubMed:11739654). Highly mobile in cells and can be relocalized through interaction with targeting subunits (PubMed:17965019). In the presence of PPP1R8 relocalizes from the nucleolus to nuclear speckles (PubMed:11739654). Shows a dynamic targeting to specific sites throughout the cell cycle (PubMed:12529430). Highly concentrated in nucleoli of interphase cells and localizes at</p>
<b>Expression :</b>	Embryonic kidney,Placenta,Skeletal muscle,

---

## Products Images



Western blot detection of PP1C in HeLa, 293T and Jurkat cell lysates using PP1C mouse mAb (1:500 diluted). Predicted band size: 38KDa. Observed band size: 38KDa.