

## AU\*Polyclonal Goat Anti Pig IgG(h+l)

|                            |                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Catalog No :</b>        | RS030436                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>Reactivity :</b>        | Pig                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Applications :</b>      | WB;IHC;IF;ELISA                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Target :</b>            | Pig IgG(h+l)                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| <b>Formulation :</b>       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Source :</b>            | Goat                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>Dilution :</b>          | WB 1:1000-10000, ELISA 1:10000-20000, IHC 1:50-100 IF 1:50-1000                                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Purification :</b>      | The antibody was affinity-purified by affinity-chromatography using antigen-specific immunogen.                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Concentration :</b>     | 2mg/mL                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <b>Storage Stability :</b> | 2°C to 8°C/1 year                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Background :</b>        | Immunoway secondary antibodies are available conjugated to enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, ImmunoHistoChemistry, ImmunoFluorescence, Flow Cytometry and ELISA. We offer high quality secondary antibodies from goat, rabbit and donkey sources for your each application. Serum adsorbed secondary antibodies are also available and are recommended for use with immunoglobulin-rich samples. |

## Products Images