

## AU\*Polyclonal Goat Anti Human IgG FC

|                            |                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Catalog No :</b>        | RS030437                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>Reactivity :</b>        | Human                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Applications :</b>      | WB;IHC;IF;ELISA                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Target :</b>            | Human IgG FC                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| <b>Formulation :</b>       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Source :</b>            | Goat                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>Dilution :</b>          | WB 1:1000-10000, ELISA 1:10000-20000, IHC 1:50-100 IF 1:50-1000                                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Purification :</b>      | The antibody was affinity-purified by affinity-chromatography using antigen-specific immunogen.                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Concentration :</b>     | 2mg/mL                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <b>Storage Stability :</b> | 2°C to 8°C/1 year                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Background :</b>        | Immunoway secondary antibodies are available conjugated to enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, ImmunoHistoChemistry, ImmunoFluorescence, Flow Cytometry and ELISA. We offer high quality secondary antibodies from goat, rabbit and donkey sources for your each application. Serum adsorbed secondary antibodies are also available and are recommended for use with immunoglobulin-rich samples. |

## Products Images