

## FITC\*Polyclonal Rabbit Anti Pig IgG(h+l)

|                            |  |
|----------------------------|--|
| <b>Catalog No :</b>        | RS030618   |
| <b>Reactivity :</b>        | Pig  |
| <b>Applications :</b>      | IF;ELISA   |
| <b>Target :</b>            | Pig IgG(h+l)   |
| <b>Formulation :</b>       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| <b>Source :</b>            | Polyclonal, Rabbit,IgG   |
| <b>Dilution :</b>          | Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are IF (1:250 - 1:2000),   |
| <b>Purification :</b>      | The antibody was affinity-purified by affinity-chromatography using antigen-specific immunogen.  |
| <b>Concentration :</b>     | 2mg/mL   |
| <b>Storage Stability :</b> | -15°C to -25°C/1 year(Do not lower than -25°C)   |
| <b>Background :</b>        | Immunoway secondary antibodies are available conjugated to enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, ImmunoHistoChemistry, ImmunoFluorescence, Flow Cytometry and ELISA. We offer high quality secondary antibodies from goat, rabbit and donkey sources for your each application. Serum adsorbed secondary antibodies are also available and are recommended for use with immunoglobulin-rich samples. |

## Products Images