

## Mouse Anti Chicken IgY (H&L) FITC

<b>Catalog No :</b>	RS2335
<b>Reactivity :</b>	Chicken
<b>Applications :</b>	IF;FCM
<b>Target :</b>	Chicken IgY (H&L) FITC
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Monoclonal, Mouse
<b>Dilution :</b>	Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are IF (1:250 - 1:2000), FCM (1:250 - 1:2000)
<b>Purification :</b>	The antibody was isolated from ascitic by immunoaffinity chromatography using antigens coupled to agarose beads.
<b>Concentration :</b>	1mg/mL
<b>Storage Stability :</b>	Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezi
<b>Background :</b>	Immunoway secondary antibodies are available conjugated to enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, ImmunoHistoChemistry, ImmunoFluorescence, Flow Cytometry and ELISA. We offer high quality secondary antibodies from goat, rabbit and donkey sources for your each application. Serum adsorbed secondary antibodies are also available and are recommended for use with immunoglobulin-rich samples.

## Products Images