

## Goat Anti Rabbit IgG (H&L)-AbFluor 488

Catalog No: RS3211

Reactivity: Rabbit

**Applications :** Elisa;IF;FCM

Target: Rabbit IgG (H&L)

**Formulation:** 1 mg/ml, liquid in 0.01M Phosphate Buffered Saline, pH 7.2, containing 1%

BSA, 50% glycerol, 0.02% Sodium Azide

Source: Goat

**Dilution:** IF (1:200 - 1:1000), FCM (1:100 - 1:1000), Elisa (Use at an assay dependent

concentration)

**Purification:** The antibody was isolated from antisera by immunoaffinity chromatography

using antigens coupled to agarose beads.

**Concentration:** 1mg/mL

Storage Stability: Stable for one year at -15°C to -25°C from date of shipment. For maximum

recovery of product, centrifuge the original vial after thawing and prior to removing

the cap. Aliquot to avoid repeated freezi


**Background:** Immunoway secondary antibodies are available conjugated to enzyme, biotin or

fluorophore for use in a variety of antibody-based applications including Western Blot, ImmunoHistoChemistry, ImmunoFluorescence, Flow Cytometry and ELISA. We offer high quality secondary antibodies from goat, rabbit and donkey sources for your each application. Serum adsorbed secondary antibodies are also


available and are recommended for use with immunoglobulin-rich samples.

## **Products Images**

1/2



Inhaled Lipid Nanoparticles Alleviate Established Pulmonary Fibrosis Small Yang Yang IF Mouse murine lung epithelial-12 (MLE-12) cell



Losartan attenuates acetic acid enema-induced visceral hypersensitivity by inhibiting the ACE1/Ang II/AT1 receptor axis in enteric glial cells EUROPEAN JOURNAL OF PHARMACOLOGY Fei Dai IF Rat colon tissue enteric glial cells (EGCs)

| 346/442 | Blue  |
|---------|---|
| 401/421 | Blue  |
| 496/519 | Green   |
| 532/553 | Yellow  |
| 555/565 | Yellow  |
| 578/603 | Red/Orange  |
| 590/617 | Red/Orange  |
| 632/647 | Red   |
| 650/665 | Red   |
| 663/690 | Near IR   |
| 679/702 | Near IR   |
| 749/775 | Near IR   |
| 784/814 | Near IR   |
|         | 401/421<br>496/519<br>532/553<br>555/565<br>578/603<br>590/617<br>632/647<br>650/665<br>663/690<br>679/702<br>749/775 |

To use the Alexa Fluors with fluorescent imagers, use a spectral line of the blue laser diode for Alexa Fluors 405, a cyan (488 nm) laser for Alexa Fluors 488, a yellow (526 nm) laser for Alexa Fluor 550 or 594, and a red (633 nm) laser for Alexa Fluor 649. The Alexa Fluor 680 and 790 fluors are compatible with laser- and filter-based infrared imaging instruments that emit in the 700 nm, and 800 nm