

SOX9 (PT0295R) PT® Rabbit mAb


Catalog No :	YM8170
Reactivity :	Human;Mouse;Rat;
Applications :	WB;IHC;IF;ELISA
Target :	Sox-9
Fields :	>>cAMP signaling pathway
Gene Name :	SOX9
Protein Name :	Transcription factor SOX-9
Human Gene Id :	6662
Human Swiss Prot No :	P48436
Specificity :	endogenous
Formulation :	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source :	Monoclonal, rabbit, IgG, Kappa
Dilution :	IHC 1:200-1:1000,WB 1:500-1:1000,IF 1:200-1:1000,ELISA 1:5000-1:20000
Purification :	Protein A
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	56kD
-	7040
Observed Band :	70kD
Background :	SRY-box 9(SOX9) Homo sapiens The protein encoded by this gene recognizes the sequence CCTTGAG along with other members of the HMG-box class DNA-binding proteins. It acts during chondrocyte differentiation and, with steroidogenic factor 1, regulates transcription of the anti-Muellerian hormone (AMH) gene.



	frequently with sex reversal. [provided by RefSeq, Jul 2008],
Function :	disease:Defects in SOX9 are the cause of campomelic dysplasia (CMD1) [MIM:114290]. CMD1 is a rare, often lethal, dominantly inherited, congenital osteochondrodysplasia, associated with male-to-female autosomal sex reversal in two-thirds of the affected karyotypic males. A disease of the newborn characterized by congenital bowing and angulation of long bones, unusually small scapulae, deformed pelvis and spine and a missing pair of ribs. Craniofacial defects such as cleft palate, micrognatia, flat face and hypertelorism are common. Various defects of the ear are often evident, affecting the cochlea, malleus incus, stapes and tympanum. Most patients die soon after birth due to respiratory distress which has been attributed to hypoplasia of the tracheobronchial cartilage and small thoracic cage.,function:Plays an important role in the normal skeletal development. May regulate the expression
Subcellular	Nucleus
Location : Expression :	Eye,PNS,Testis,

Deficiencies lead to the skeletal malformation syndrome campomelic dysplasia,

Products Images



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-SOX9 (PT0295R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: HepG2 Lane 2: U-14 Lane 3: SW480 Predicted band size: 56kDa Observed band size: 70kDa