

JAK1 pro	tein
----------	------

Catalog No: YD0055

Reactivity: Human

Applications: WB;SDS-PAGE

Gene Name: JAK1

Protein Name: JAK1 protein

Sequence: Amino acid: 450-618, with his-MBP tag.

P23458

P52332

Human Gene ld: 3716

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Formulation : Liquid in PBS

Source : E.coli

Dilution : WB 1:500-2000

Concentration: SDS-PAGE >90%

Storage Stability: -20°C/6 month,-80°C for long storage

Background : catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine

phosphate.,domain:Possesses two phosphotransferase domains. The second one probably contains the catalytic domain (By similarity), while the presence of slight differences suggest a different role for domain 1.,domain:The FERM domain mediates interaction with JAKMIP1.,function:Tyrosine kinase of the non-receptor type, involved in the IFN-alpha/beta/gamma signal pathway. Kinase partner for the interleukin (IL)-2 receptor.,sequence caution:Translation N-terminally extended.,similarity:Belongs to the protein kinase superfamily. Tyr

protein kinase family. JAK subfamily., similarity: Contains 1 FERM

domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 SH2

domain., subcellular location: Wholly intracellular, possibly membrane



associated.,subunit:Interacts with IL31RA, JAKMIP1 and SHB.,tissue specificity:Expressed at higher levels in primary colon tumors than in normal colon tissue. The expression level in metastatic colon tumors is comparable to the expression level in normal colon tissue.,

Function:

protein amino acid phosphorylation, phosphorus metabolic process, phosphate metabolic process, cell surface receptor linked signal transduction, enzyme linked receptor protein signaling pathway, intracellular signaling cascade, protein kinase cascade, phosphorylation, peptidyl-tyrosine phosphorylation, peptidyl-tyrosine modification, cytokine-mediated signaling pathway, response to antibiotic,

Subcellular Location:

Endomembrane system; Peripheral membrane protein. Wholly intracellular, possibly membrane associated.

Expression:

Expressed at higher levels in primary colon tumors than in normal colon tissue. The expression level in metastatic colon tumors is comparable to the expression level in normal colon tissue.

Products Images

