

Recombinant Human MUC1 protein ,N- His Tag

Catalog No :	YD2288
Reactivity :	Human
Purity :	>90% as determined by SDS-PAGE
In Vivo :	Please contact with the lab for this information.
Protein Name :	Mucin-1 (MUC-1) (Breast carcinoma-associated antigen DF3) (Carcinoma-associated mucin) (Episialin) (H23AG) (Krebs von den Lungen-6) (KL-6) (PEMT) (Peanut-reactive urinary mucin) (PUM) (Polymorphic epi
Sequence :	A DNA sequence encoding the Human MUC1 (Ser24-Pro146 [?]) was fused with the N-terminal His Tag
Human Swiss Prot No :	P15941
Formulation :	Supplied as solution form in PBS pH 7.5 ./ Lyophilized from PBS pH 7.5 .
Source :	E. coli
Dilution :	Reconstitute in sterile water for a stock solution
Concentration :	>90% as determined by SDS-PAGE
Storage Stability :	Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 2 to 8 °C for one week . Store at -20 to -80 °C for twelve months from the date of receipt.
Subcellular Location :	Apical cell membrane ; Single-pass type I membrane protein . Exclusively located in the apical domain of the plasma membrane of highly polarized epithelial cells. After endocytosis, internalized and recycled to the cell membrane. Located to microvilli and to the tips of long filopodial protrusions.; [Isoform 5]: Secreted.; [Isoform Y]: Secreted.; [Isoform 9]: Secreted.; [Mucin-1 subunit beta]: Cell membrane. Cytoplasm. Nucleus. On EGF and PDGFRB stimulation, transported to the nucleus through interaction with CTNNB1, a process which is stimulated by phosphorylation. On HRG stimulation, colocalizes with JUP/gamma-catenin at the nucleus.

Expression : Expressed on the apical surface of epithelial cells, especially of airway passages, breast and uterus. Also expressed in activated and unactivated T-cells. Overexpressed in epithelial tumors, such as breast or ovarian cancer and also in non-epithelial tumor cells. Isoform Y is expressed in tumor cells only.

Products Images