

**Oct-4 (PTR1369) mouse mAb**

<b>Catalog No :</b>	YM4770
<b>Reactivity :</b>	Human;
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	OCT-4
<b>Fields :</b>	>>Signaling pathways regulating pluripotency of stem cells
<b>Gene Name :</b>	POU5F1
<b>Protein Name :</b>	POU domain class 5 transcription factor 1
<b>Human Gene Id :</b>	5460
<b>Human Swiss Prot No :</b>	Q01860
<b>Immunogen :</b>	Synthesized peptide derived from human Oct-4 AA range:50-150
<b>Specificity :</b>	This antibody detects endogenous levels of 4-Oct protein.
<b>Formulation :</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
<b>Source :</b>	Monoclonal, Mouse,IgG2b,kappa
<b>Dilution :</b>	WB 1:500 - 1:2000. ELISA: 1:1000-5000.
<b>Purification :</b>	The antibody was affinity-purified from ascites by affinity-chromatography using specific immunogen.
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Molecularweight :</b>	39kD
<b>Observed Band :</b>	40kD

**Background :**

This gene encodes a transcription factor containing a POU homeodomain that plays a key role in embryonic development and stem cell pluripotency. Aberrant expression of this gene in adult tissues is associated with tumorigenesis. This gene can participate in a translocation with the Ewing's sarcoma gene on chromosome 21, which also leads to tumor formation. Alternative splicing, as well as usage of alternative AUG and non-AUG translation initiation codons, results in multiple isoforms. One of the AUG start codons is polymorphic in human populations. Related pseudogenes have been identified on chromosomes 1, 3, 8, 10, and 12. [provided by RefSeq, Oct 2013],

**Function :**

function:Transcription factor that binds to the octamer motif (5'-ATTTGCAT-3'). Forms a trimeric complex with SOX2 on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206. Critical for early embryogenesis and for embryonic stem cell pluripotency.,miscellaneous:Several pseudogenes of POU5F1 have been described on chromosomes 1, 3, 8, 10 and 12. 2 of them, localized in chromosomes 8 and 10, are transcribed in cancer tissues but not in normal ones and may be involved in the regulation of POU5F1 gene activity in carcinogenesis.,online information:Oct-4 entry,PTM:Sumoylation enhances the protein stability, DNA binding and transactivation activity. Sumoylation is required for enhanced YES1 expression.,similarity:Belongs to the POU transcription factor family. Class-5 subfamily.,similarity:Contains 1 homeobox DNA-binding doma


**Subcellular Location :**

Cytoplasm. Nucleus. Note=Expressed in a diffuse and slightly punctuate pattern. Colocalizes with MAPK8 and MAPK9 in the nucleus.

**Expression :**

Expressed in developing brain. Highest levels found in specific cell layers of the cortex, the olfactory bulb, the hippocampus and the cerebellum. Low levels of expression in adult tissues.

## Products Images



Whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with anti Oct-4 (PTR1369) antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: NCCIT Predicted band size: 39kDa Observed band size: 40kDa