

CD14-FC recombinant protein

Catalog No :	YD3062
Reactivity :	Human;
Purity :	>90% as determined by SDS-PAGE
Gene Name :	CD14
Protein Name :	Monocyte differentiation antigen CD14 (Myeloid cell-specific leucine-rich glycoprotein) (CD antigen CD14) [Cleaved into: Monocyte differentiation antigen CD14, urinary form; Monocyte differentiation a
Sequence :	Amino acid:19-345,with FC tag.
Human Gene Id :	929
Human Swiss Prot No :	P08571
Formulation :	Phosphate-buffered solution
Source :	Mammalian cells
Storage Stability :	-15°C to -25°C/1 year(Avoid freeze / thaw cycles)
Function :	Coreceptor for bacterial lipopolysaccharide (PubMed:1698311, PubMed:23264655). In concert with LBP, binds to monomeric lipopolysaccharide and delivers it to the LY96/TLR4 complex, thereby mediating the innate immune response to bacterial lipopolysaccharide (LPS) (PubMed:20133493, PubMed:22265692, PubMed:23264655). Acts via MyD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed:8612135). Acts as a coreceptor for TLR2:TLR6 heterodimer in response to diacylated lipopeptides and for TLR2:TLR1 heterodimer in response to triacylated lipopeptides, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway (PubMed:16880211). Binds electronegative LDL (LDL(-)) and mediates the cytokine release induced by LDL(-) (PubMed:23880187).
Subcellular Location :	Cell membrane ; Lipid-anchor, GPI-anchor . Secreted . Membrane raft . Golgi apparatus . Note=Secreted forms may arise by cleavage of the GPI anchor. .

Expression : Detected on macrophages (at protein level) (PubMed:1698311). Expressed strongly on the surface of monocytes and weakly on the surface of granulocytes; also expressed by most tissue macrophages.

Products Images