

RAGE rabbit-FC recombinant protein

YD3132 Catalog No:

Reactivity: Human;

>90% as determined by SDS-PAGE **Purity:**

Gene Name: RAGE

Protein Name: Advanced glycosylation end product-specific receptor

Amino acid:26-340, with rabbit FC tag. Sequence:

Human Gene Id: 177

Human Swiss Prot

No:

Formulation: Phosphate-buffered solution

Q15109

Source: Mammalian cells

Storage Stability: -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

The advanced glycosylation end product (AGE) receptor encoded by this gene **Background:**

> is a member of the immunoglobulin superfamily of cell surface receptors. It is a multiligand receptor, and besides AGE, interacts with other molecules implicated in homeostasis, development, and inflammation, and certain diseases, such as diabetes and Alzheimer's disease. Many alternatively spliced transcript variants encoding different isoforms, as well as non-protein-coding variants, have been described for this gene (PMID:18089847). [provided by RefSeg, May 2011],

Function: function: Mediates interactions of advanced glycosylation end products (AGE).

> These are nonenzymatically glycosylated proteins which accumulate in vascular tissue in aging and at an accelerated rate in diabetes. Receptor for amyloid beta

peptide., similarity: Contains 1 Ig-like V-type (immunoglobulin-like) domain., similarity: Contains 2 Ig-like C2-type (immunoglobulin-like)

domains., tissue specificity: Endothelial cells.,

Subcellular [Isoform 1]: Cell membrane; Single-pass type I membrane protein.; [Isoform 2]: Location:

Secreted.; [Isoform 10]: Cell membrane; Single-pass type I membrane protein.



Expression : Endothelial cells.

Products Images

2/2