

NOX4 Polyclonal Antibody

Catalog No :	YN2975
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	NOX4
Fields :	>>AGE-RAGE signaling pathway in diabetic complications;>>Alcoholic liver disease;>>Alzheimer disease;>>Pathways of neurodegeneration - multiple diseases;>>Chemical carcinogenesis - reactive oxygen species
Gene Name :	NOX4 RENOX
Protein Name :	NADPH oxidase 4 (EC 1.6.3.-) (Kidney oxidase-1) (KOX-1) (Kidney superoxide-producing NADPH oxidase) (Renal NAD(P)H-oxidase)
Human Gene Id :	50507
Human Swiss Prot No :	Q9NPH5
Mouse Swiss Prot No :	Q9JHI8
Rat Swiss Prot No :	Q924V1
Immunogen :	Synthesized peptide derived from part region of human protein AA range:520-578
Specificity :	NOX4 Polyclonal Antibody detects endogenous levels of protein.
Formulation :	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000 ELISA 1:5000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 63kD

Background : This gene encodes a member of the NOX-family of enzymes that functions as the catalytic subunit the NADPH oxidase complex. The encoded protein is localized to non-phagocytic cells where it acts as an oxygen sensor and catalyzes the reduction of molecular oxygen to various reactive oxygen species (ROS). The ROS generated by this protein have been implicated in numerous biological functions including signal transduction, cell differentiation and tumor cell growth. A pseudogene has been identified on the other arm of chromosome 11. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Jan 2009],

Function : developmental stage:Expressed in fetal kidney and fetal liver.,enzyme regulation:Inhibited by plumbagin (By similarity). Activated by phorbol 12-myristate 13-acetate (PMA). Activated by insulin. Inhibited by diphenylene iodonium.,function:Constitutive NADPH oxidase which generates superoxide intracellularly upon formation of a complex with CYBA/p22phox. Regulates signaling cascades probably through phosphatases inhibition. May function as an oxygen sensor regulating the KCNK3/TASK-1 potassium channel and HIF1A activity. May regulate insulin signaling cascade. May play a role in apoptosis, bone resorption and lipopolysaccharide-mediated activation of NFkB. Isoform 3 is not functional. Isoform 4 displays an increased activity while isoform 5 and isoform 6 display reduced activity. May produce superoxide in the nucleus and play a role in regulating gene expression upon cell stimulation.,induc

Subcellular Location : Endoplasmic reticulum membrane ; Multi-pass membrane protein. Cell membrane ; Multi-pass membrane protein . Cell junction, focal adhesion . May localize to plasma membrane and focal adhesions. According to PubMed:15927447, may also localize to the nucleus.; [Isoform 4]: Nucleus . Nucleus, nucleolus .

Expression : Expressed by distal tubular cells in kidney cortex and in endothelial cells (at protein level). Widely expressed. Strongly expressed in kidney and to a lower extent in heart, adipocytes, hepatoma, endothelial cells, skeletal muscle, brain, several brain tumor cell lines and airway epithelial cells.

Products Images