

Sumo 1 (PT0513R) PT® Rabbit mAb

Catalog No :	YM8339
Reactivity :	Human; Mouse; Rat;
Applications :	WB;IHC;IF;IP;ELISA
Target :	SUMO-1
Fields :	>>Nucleocytoplasmic transport;>>Fluid shear stress and atherosclerosis
Gene Name :	SUMO1
Protein Name :	Small ubiquitin-related modifier 1
Human Gene Id :	7341
Human Swiss Prot No :	P63165
Mouse Gene Id :	22218
Mouse Swiss Prot No :	P63166
Rat Gene Id :	301442
Rat Swiss Prot No :	Q5I0H3
Specificity :	endogenous
Formulation :	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source :	Monoclonal, rabbit, IgG, Kappa
Dilution :	IHC 1:2000-1:10000;WB 1:2000-1:10000;IF 1:200-1:1000;ELISA 1:5000-1:20000;IP 1:50-1:200;
Purification :	Protein A

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight : 12kD

Observed Band : 80kD


Background : This gene encodes a protein that is a member of the SUMO (small ubiquitin-like modifier) protein family. It functions in a manner similar to ubiquitin in that it is bound to target proteins as part of a post-translational modification system. However, unlike ubiquitin which targets proteins for degradation, this protein is involved in a variety of cellular processes, such as nuclear transport, transcriptional regulation, apoptosis, and protein stability. It is not active until the last four amino acids of the carboxy-terminus have been cleaved off. Several pseudogenes have been reported for this gene. Alternate transcriptional splice variants encoding different isoforms have been characterized. [provided by RefSeq, Jul 2008],

Function : caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,function:Ubiquitin-like protein which can be covalently attached to target lysines as a monomer. Does not seem to be involved in protein degradation and may function as an antagonist of ubiquitin in the degradation process. Plays a role in a number of cellular processes such as nuclear transport, DNA replication and repair, mitosis and signal transduction. Involved in targeting RANGAP1 to the nuclear pore complex protein RANBP2. Covalent attachment to its substrates requires prior activation by the E1 complex SAE1-SAE2 and linkage to the E2 enzyme UBE2I, and can be promoted by an E3 ligase such as PIAS1-4, RANBP2 or CBX4.,online information:SUMO protein entry,PTM: Cleavage of precursor form by SENP1 or SENP2 is necessary for function.,similarity:Belongs to t


Subcellular Location : Cytoplasm, Nucleus

Expression : Brain,Colon adenocarcinoma,Epithelium,Placenta,

Products Images



Human lung carcinoma was stained with anti-Sumo 1 (PT0513R) rabbit antibody



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-Sumo 1 (PT0513R) antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: Jurkat Lane 2: HeLa Lane 3: PC-12 Lane 4: NIH-3T3 Predicted band size: 12kDa Observed band size: 80kDa