

INSRR Polyclonal Antibody

Catalog No: YN2275

Reactivity: Human;Rat;Mouse

Applications: WB;ELISA

Target: INSRR

Fields: >>Regulation of actin cytoskeleton;>>Prostate cancer

Gene Name: INSRR IRR

Protein Name: Insulin receptor-related protein (IRR) (EC 2.7.10.1) (IR-related receptor)

[Cleaved into: Insulin receptor-related protein alpha chain; Insulin receptor-related

protein beta chain]

Q9WTL4

Human Gene Id: 3645

Human Swiss Prot P14616

No:

Mouse Swiss Prot

No:

Rat Swiss Prot No: Q64716

Immunogen: Synthesized peptide derived from human protein. at AA range: 80-160

Specificity: INSRR Polyclonal Antibody detects endogenous levels of protein.

Formulation : Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500-2000 ELISA 1:5000-20000

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

1/2



Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 142kD

Cell Pathway: Regulates Actin and Cytoskeleton; Prostate cancer;

Background: catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine

phosphate.,function:This receptor probably binds an insulin related protein and has a tyrosine-protein kinase activity. It phosphorylates the insulin receptor substrates IRS-1 and IRS-2.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily.,similarity:Contains 1 protein

kinase domain., similarity: Contains 3 fibronectin type-III

domains., subunit: Probable tetramer of 2 alpha and 2 beta chains linked by disulfide bonds. The alpha chains contribute to the formation of the ligand-binding

domain, while the beta chains carry the kinase domain.,

Function: catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine

phosphate.,function:This receptor probably binds an insulin related protein and has a tyrosine-protein kinase activity. It phosphorylates the insulin receptor substrates IRS-1 and IRS-2.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily.,similarity:Contains 1 protein

kinase domain., similarity: Contains 3 fibronectin type-III

domains., subunit: Probable tetramer of 2 alpha and 2 beta chains linked by disulfide bonds. The alpha chains contribute to the formation of the ligand-binding

domain, while the beta chains carry the kinase domain.

Subcellular Location:

Membrane; Single-pass type I membrane protein.

Expression:

Kidney,

Products Images