

## Tissue Inhibitor of Metalloproteinases 1(TIMP1) (ABT200R) rabbit mAb

<b>Catalog No :</b>	YM7224
<b>Reactivity :</b>	Human;
<b>Applications :</b>	WB; ELISA
<b>Target :</b>	TIMP1
<b>Fields :</b>	>>HIF-1 signaling pathway
<b>Gene Name :</b>	TIMP1
<b>Protein Name :</b>	Tissue Inhibitor of Metalloproteinases 1(TIMP1)
<b>Human Gene Id :</b>	7076
<b>Human Swiss Prot No :</b>	P01033
<b>Immunogen :</b>	Synthesized peptide derived from human Tissue Inhibitor of Metalloproteinases 1(TIMP1) AA range:50-150
<b>Specificity :</b>	This antibody detects endogenous levels of TIMP1
<b>Formulation :</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
<b>Source :</b>	Monoclonal, Rabbit IgG1, Kappa
<b>Dilution :</b>	WB 1:500-1000, ELISA 1:5000-20000
<b>Purification :</b>	Recombinant Expression and Affinity purified
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Molecularweight :</b>	23kD
<b>Background :</b>	This gene belongs to the TIMP gene family. The proteins encoded by this gene family are natural inhibitors of the matrix metalloproteinases (MMPs), a group of peptidases involved in degradation of the extracellular matrix. In addition to its

inhibitory role against most of the known MMPs, the encoded protein is able to promote cell proliferation in a wide range of cell types, and may also have an anti-apoptotic function. Transcription of this gene is highly inducible in response to many cytokines and hormones. In addition, the expression from some but not all inactive X chromosomes suggests that this gene inactivation is polymorphic in human females. This gene is located within intron 6 of the synapsin I gene and is transcribed in the opposite direction. [provided by RefSeq, Jul 2008],

---

**Function :**

function:Complexes with metalloproteinases (such as collagenases) and irreversibly inactivates them. Also mediates erythropoiesis in vitro; but, unlike IL-3, it is species-specific, stimulating the growth and differentiation of only human and murine erythroid progenitors. Known to act on MMP-1, MMP-2, MMP-3, MMP-7, MMP-8, MMP-9, MMP-10, MMP-11, MMP-12, MMP-13 and MMP-16. Does not act on MMP-14.,PTM:The activity of TIMP1 is dependent on the presence of disulfide bonds.,similarity:Belongs to the protease inhibitor I35 (TIMP) family.,similarity:Contains 1 NTR domain.,

---

**Subcellular Location :**

Secreted .

---

**Expression :**

Detected in rheumatoid synovial fluid (at protein level).

---

## Products Images