

**PIK3CA H1047R (PT0281R) rabbit mAb**

<b>Catalog No :</b>	YM7195
<b>Reactivity :</b>	Human;
<b>Applications :</b>	IHC;WB; ELISA
<b>Target :</b>	PIK3CA H1047R
<b>Fields :</b>	>>Inositol phosphate metabolism;>>Metabolic pathways;>>EGFR tyrosine kinase inhibitor resistance;>>Endocrine resistance;>>Platinum drug resistance;>>ErbB signaling pathway;>>Ras signaling pathway;>>Rap1 signaling pathway;>>cAMP signaling pathway;>>Chemokine signaling pathway;>>HIF-1 signaling pathway;>>FoxO signaling pathway;>>Phosphatidylinositol signaling system;>>Sphingolipid signaling pathway;>>Phospholipase D signaling pathway;>>Autophagy - animal;>>mTOR signaling pathway;>>PI3K-Akt signaling pathway;>>AMPK signaling pathway;>>Apoptosis;>>Longevity regulating pathway;>>Longevity regulating pathway - multiple species;>>Cellular senescence;>>Axon guidance;>>VEGF signaling pathway;>>Osteoclast differentiation;>>Focal adhesion;>>Signaling pathways regulating pluripotency of stem cells;>>Platelet activation;>>Neutrophil extracellular trap formation;>>Toll-like receptor signaling pathway;>>C-type lectin receptor signaling pathway;>>JAK-STAT signaling pathway;>>Natural killer cell mediat
<b>Gene Name :</b>	PIK3CA
<b>Protein Name :</b>	5-bisphosphate 3-kinase 110 kDa catalytic subunit alpha;5-bisphosphate 3-kinase catalytic subunit alpha isoform;caPI3K;CLOVE;CWS5;MCAP;MCM;MC MTC;MGC142161;MGC142163;p110 alpha;p110alpha;Phosphatidylin
<b>Human Swiss Prot No :</b>	P42336
<b>Mouse Swiss Prot No :</b>	P42337
<b>Immunogen :</b>	Synthesized peptide derived from human PIK3CA H1047R AA range:1000-1068
<b>Specificity :</b>	This antibody detects endogenous levels of PIK3CA H1047R
<b>Formulation :</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA



Best Tools for immunology Research

<b>Source :</b>	Monoclonal, Rabbit IgG1, Kappa
<b>Dilution :</b>	IHC 1:100-500, WB 1:500-1000, ELISA 1:5000-20000
<b>Purification :</b>	Recombinant Expression and Affinity purified
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Molecularweight :</b>	117kD
<b>Background :</b>	Phosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. The protein encoded by this gene represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has been found to be oncogenic and has been implicated in cervical cancers. A pseudogene of this gene has been defined on chromosome 22. [provided by RefSeq, Apr 2016],
<b>Function :</b>	catalytic activity:ATP + 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate = ADP + 1-phosphatidyl-1D-myo-inositol 3,4,5-trisphosphate.,disease:Defects in PIK3CA are associated with breast cancer [MIM:114480].,disease:Defects in PIK3CA are associated with colorectal cancer (CRC) [MIM:114500].,disease:Defects in PIK3CA are associated with ovarian cancer [MIM:167000]. Ovarian cancer is the leading cause of death from gynecologic malignancy. It is characterized by advanced presentation with loco-regional dissemination in the peritoneal cavity and the rare incidence of visceral metastases. These typical features relate to the biology of the disease, which is a principal determinant of outcome.,disease:Defects in PIK3CA may underlie hepatocellular carcinoma (HCC) [MIM:114550].,disease:PI3KCA mutations affecting exons 9 and 20 display gender-and tissue-specific patterns, thus suggesting that the
<b>Subcellular Location :</b>	Cytoplasmic
<b>Expression :</b>	Gastric adenocarcinoma with PIK3CA H1047R protein expression

## Products Images