

Claudin 4 (ABT188R) rabbit mAb

Catalog No: YM7088

Reactivity: Human;

Applications: WB;IHC; ELISA

Target: Claudin-4

Fields: >>Cell adhesion molecules;>>Tight junction;>>Leukocyte transendothelial

migration;>>Pathogenic Escherichia coli infection;>>Hepatitis C

Gene Name: CLDN4

Protein Name: Claudin-4 (Clostridium perfringens enterotoxin receptor) (CPE-R) (CPE-

receptor) (Williams-Beuren syndrome chromosomal region 8 protein)

Human Gene Id: 1364

Human Swiss Prot

No:

Immunogen: Synthesized peptide derived from human Claudin 4 AA range:150-209

Specificity: This antibody detects endogenous levels of Claudin-4

Formulation: PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

Source: Monoclonal, Rabbit IgG1, Kappa

O14493

Dilution: IHC 1:100-500, WB 1:500-1000, ELISA 1:5000-20000

Purification: Recombinant Expression and Affinity purified

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 22kD

Background: The protein encoded by this intronless gene belongs to the claudin family.

Claudins are integral membrane proteins that are components of the epithelial cell



tight junctions, which regulate movement of solutes and ions through the paracellular space. This protein is a high-affinity receptor for Clostridium perfringens enterotoxin (CPE) and may play a role in internal organ development and function during pre- and postnatal life. This gene is deleted in Williams-Beuren syndrome, a neurodevelopmental disorder affecting multiple systems. [provided by RefSeq, Sep 2013],

Function: disease:Haploinsufficiency of CLDN4 may be the cause of certain

cardiovascular and musculo-skeletal abnormalities observed in Williams-Beuren syndrome (WBS), a rare developmental disorder. It is a contiguous gene deletion syndrome involving genes from chromosome band 7q11.23., function: Plays a

major role in tight junction-specific obliteration of the intercellular

space., similarity: Belongs to the claudin family., subunit: Directly interacts with

TJP1/ZO-1, TJP2/ZO-2 and TJP3/ZO-3.,

Subcellular Location:

Cell junction, tight junction. Cell membrane; Multi-pass membrane protein.

CLDN4 is required for tight junction localization in the kidney. .

Expression: Colon, Fetal brain, Trachea,

Products Images