

DUSP6 Polyclonal Antibody

Catalog No :	YN5617
Reactivity :	Human;Rat;Mouse
Applications :	WB
Target :	DUSP6
Fields :	>>MAPK signaling pathway;>>Transcriptional misregulation in cancer;>>Acute myeloid leukemia
Gene Name :	DUSP6
Protein Name :	Dual specificity protein phosphatase 6 (EC 3.1.3.16) (EC 3.1.3.48) (Dual specificity protein phosphatase PYST1) (Mitogen-activated protein kinase phosphatase 3) (MAP kinase phosphatase 3) (MKP-3)
Human Gene Id :	1848
Human Swiss Prot No :	Q16828
Mouse Swiss Prot No :	Q9DBB1
Rat Swiss Prot No :	Q64346
Immunogen :	Synthetic Peptide of DUSP6
Specificity :	DUSP6 Rabbit Polyclonal detects endogenous levels of DUSP6
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:1000-3000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration :	<u>1 mg/ml</u>
Storage Stability :	<u>-15°C to -25°C/1 year(Do not lower than -25°C)</u>
Observed Band :	<u>42,44kD</u>
Cell Pathway :	<u>MAPK_ERK_Growth;MAPK_G_Protein;</u>
Background :	<p>The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product inactivates ERK2, is expressed in a variety of tissues with the highest levels in heart and pancreas, and unlike most other members of this family, is localized in the cytoplasm. Mutations in t</p>
Function :	<p>catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate.,catalytic activity:Protein tyrosine phosphate + H(2)O = protein tyrosine + phosphate.,function:Inactivates MAP kinases. Has a specificity for the ERK family.,similarity:Belongs to the protein-tyrosine phosphatase family. Non-receptor class dual specificity subfamily.,similarity:Contains 1 rhodanese domain.,similarity:Contains 1 tyrosine-protein phosphatase domain.,</p>
Subcellular Location :	<u>Cytoplasm .</u>
Expression :	<u>Expressed in keratinocytes (at protein level).</u>

Products Images

Western blot analysis of 1) HepG2, 2)3T3, 3) Rat Heart Tissue with DUSP6 Rabbit pAb diluted at 1:3,000.

