

Cytokeratin 8 (CK8) (ABT170R) rabbit mAb (Ready to Use)

Catalog No :	YM7263R
Reactivity :	Human;
Applications :	IHC
Target :	Cytokeratin 8
Gene Name :	KRT8 CYK8
Protein Name :	CARD2;CK 8;CK-8;CK8;CYK8;CYKER;Cytokeratin endo A;Cytokeratin-8;DreK8;EndoA;K2C8;K2C8_HUMAN;K8;Keratin 8;Keratin type II cytoskeletal 8;Keratin, type II cytoskeletal 8;Keratin-8;KO;Krt 2.8;KRT8;MGC118
Human Swiss Prot No :	P05787
Mouse Swiss Prot No :	P11679
Rat Swiss Prot No :	Q10758
Immunogen :	Synthesized peptide derived from human Cytokeratin 8 AA range:100-200
Specificity :	This antibody detects endogenous levels of Cytokeratin 8
Formulation :	The prediluted ready-to-use antibody is diluted in phosphate buffer saline containing stabilizing protein and 0.05% Proclin 300
Source :	Monoclonal, Rabbit IgG1, Kappa
Dilution :	Ready to use for IHC
Purification :	Recombinant Expression and Affinity purified
Storage Stability :	2°C to 8°C/1 year
Molecularweight :	53kD
Background :	keratin 8(KRT8) Homo sapiens This gene is a member of the type II keratin

family clustered on the long arm of chromosome 12. Type I and type II keratins heteropolymerize to form intermediate-sized filaments in the cytoplasm of epithelial cells. The product of this gene typically dimerizes with keratin 18 to form an intermediate filament in simple single-layered epithelial cells. This protein plays a role in maintaining cellular structural integrity and also functions in signal transduction and cellular differentiation. Mutations in this gene cause cryptogenic cirrhosis. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jan 2012],

Function :

disease:Defects in KRT8 are a cause of cryptogenic cirrhosis [MIM:215600].,function:Together with KRT19, helps to link the contractile apparatus to dystrophin at the costameres of striated muscle.,miscellaneous:There are two types of cytoskeletal and microfibrillar keratin: I (acidic; 40-55 kDa) and II (neutral to basic; 56-70 kDa).,PTM:O-glycosylated at multiple sites; glycans consist of single N-acetylglucosamine residues.,PTM:Phosphorylation on serine residues is enhanced during EGF stimulation and mitosis. Ser-74 phosphorylation plays an important role in keratin filament reorganization.,similarity:Belongs to the intermediate filament family.,subunit:Heterotetramer of two type I and two type II keratins. keratin-8 associates with keratin-18. Associates with KRT20. Interacts with HCV core protein and PNN. When associated with KRT19, interacts with DMD. Interacts with TCHP.,tissue spec


Subcellular Location :

Cytoplasmic, Membranous


Expression :

Liver/ Tonsil

Products Images



Immunohistochemical analysis of paraffin-embedded human Colon carcinoma. 1, Antibody was incubated at 4° overnight. 2, TRIS-EDTA of pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded human Prostate. 1, Antibody was incubated at 4 ° overnight. 2, TRIS-EDTA of pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 30min).