

Beclin 1 mouse mAb

CatalogNo: YM1326 Orthogonal Validated 💽

Comparable Abs C Protein Overexpression P

Key Features

Host Species

Mouse

Reactivity

Human, Mouse, Rat, Monkey, Hamster

Applications

WB,IP

MW

55kD (Observed)

Recommended Dilution Ratios

WB 1:1000 IP 1:50-200

Storage

Storage*

-15°C to -25°C/1 year(Do not lower than -25°C)

I Basic Information

Clonality

Monoclonal

Immunogen Information

Immunogen Purified recombinant human Beclin 1 protein fragments expressed in E.coli.

Specificity This antibody detects endogenous levels of Beclin 1 and does not cross-react with

related proteins.

| Target Information

Gene name

becn1

Protein Name Organism Gene ID UniProt ID Human 8678; Q14457; Mouse 088597;

Cellular Localization


Cytoplasm . Golgi apparatus, trans-Golgi network membrane ; Peripheral membrane protein . Endosome membrane ; Peripheral membrane protein . Endoplasmic reticulum membrane ; Peripheral membrane protein . Mitochondrion membrane ; Peripheral membrane protein . Endosome . Cytoplasmic vesicle, autophagosome . Interaction with ATG14 promotes translocation to autophagosomes. Expressed in dendrites and cell bodies of cerebellar Purkinje cells (By similarity). .; [Beclin-1-C 35 kDa]: Mitochondrion . Nucleus . Cytoplasm .; [Beclin-1-C 37 kDa]: Mitochondrion .

Tissue specificity Ubiquitous.


Function

Function:Plays a central role in autophagy (By similarity). May play a role in antiviral host defense. Protects against infection by a neurovirulent strain of Sindbis virus., similarity:Belongs to the beclin family., subcellular location:Expressed in dendrites and cell bodies of cerebellar Purkinje cells., subunit:Interacts with GOPC and GRID2. Interacts with AMBRA1. Probably forms a complex with AMBRA1 and PIK3C3 (By similarity). Interacts with BCL2 and BCL2L1., tissue specificity:Ubiquitous.,


Validation Data



Deng, Z., Sun, M., Wu, J. et al. SIRT1 attenuates sepsis-induced acute kidney injury via Beclin1 deacetylation-mediated autophagy activation. Cell Death Dis 12, 217 (2021).



Deng, Z., Sun, M., Wu, J. et al. SIRT1 attenuates sepsis-induced acute kidney injury via Beclin1 deacetylation-mediated autophagy activation. Cell Death Dis 12, 217 (2021).



Western blot detection of Beclin 1 in Hela,3T3,C6,COS7,CHO-K1,Rat brain and Jurkat cell lysates using Beclin 1 mouse mAb (1:1000 diluted).Predicted band size:52KDa.Observed band size:55KDa.

Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

408-747-0189 (USA) 400-8787-807(China) Telephone:

http://www.immunoway.com Website:

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

Beclin 1 mouse mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents